



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

### Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

### About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>



## NEW WORK FOR UPPER AND MIDDLE-CLASS SCHOOLS.

By E. T. STEVENS, A.K.C., London.

AND

CHARLES HOLE, F.R.G.S.

*Editors of 'The Grade Lesson Books,' 'The Advanced Lesson Books,' &c.*

It has been suggested by numerous teachers of upper and middle-class schools that the carefully graduated system of teaching to read and spell adopted in the 'Grade Lesson Books,' and highly approved by the teachers of elementary schools under Government inspection, would find equal favour among teachers of schools of a higher class, as the principles of education are the same for the rich as for the poor.

Certain features of the 'Grade Lesson Books,' however (as, for instance), the writing and arithmetical exercises, which make them peculiarly useful in elementary schools for the poor, render them at the same time unsuited to others. In accordance with these suggestions, the Editors have resolved to publish a new series of reading and spelling books especially designed for upper and middle-class schools, to be called

## THE COMPLETE READER,

In Four Books: Being a carefully graduated System of teaching to Read and Spell by means of Attractive and Instructive Lessons, designed for Upper and Middle-Class Schools. By E. T. STEVENS, A.K.C. London, Head Master, Cathedral School, Bristol, and CHARLES HOLE, F.R.G.S. Head Master, Loughborough Collegiate School, Brixton; late Head Master S. Thomas's Collegiate School, Colombo, Ceylon; Editors of 'The Grade Lesson Books.'

**Book I.—THE PRIMARY READER;** a Course of Progressive Reading and Spelling Lessons in Monosyllables and Dissyllables (Prose and Poetry).

**Book II.—THE INTERMEDIATE READER;** a Course of Progressive Reading and Spelling Lessons (Prose and Poetry).

**Book III.—THE EXEMPLAR OF STYLE;** a Course of Reading Lessons, with Definitions, &c. consisting of careful Selections from the English Classics and Current Literature (Prose and Poetry).

**Book IV.—THE SENIOR-CLASS READER;** a Course of Lessons, with Definitions, &c. in History, Geography, Literature (Prose and Poetry), and Science, selected from the Works of the most Eminent Writers.

The Primary Reader and the Intermediate Reader have been produced to enable the pupil, by systematic and careful gradation, to read with ease, fluency, and expression; and to acquire from the first a habit of correct Spelling.

The Exemplar of Style, while attending to these points, is designed for the cultivation of the pupil's literary taste.

The Senior-Class Reader is intended to continue this training, presenting, at the same time, more solid and direct instruction in various important branches of knowledge.

---

*London:* LONGMANS, GREEN, and CO. Paternoster Row.

---





AN  
EASY INTRODUCTION  
TO THE HIGHER TREATISES ON THE  
CONIC SECTIONS.

BY THE  
REV. JOHN HUNTER, M.A.



LONDON:  
LONGMANS, GREEN, AND CO.  
1866.

*183. g. 16.*

LONDON  
PRINTED BY SPOTTISWOODE AND CO.  
NEW-STREET SQUARE

## PREFACE.

---

THE PRESENT INTRODUCTION has resulted not from theorizing on what may be expedient, but from long experience of what is needful. The author has, from time to time, had applications for assistance in the study of such treatises on the Conic Sections as those of Salmon, Hymers, Puckle, and Todhunter. He has accordingly ascertained the nature of the difficulties that perplex and discourage many readers of those treatises, and has endeavoured in this publication to provide a first course of lessons and exercises which, he thinks, cannot fail of qualifying the young student for reading with intelligence and profit the masterly works referred to. Todhunter's book is especially recommended as the most appropriate sequel to this Introduction.

It should be remarked that the name *Conic Sections* chiefly refers to the Ellipse, Hyperbola, and Parabola, as curves that can be obtained by means of a right cone variously intersected by a plane.

LONDON: *October*, 1866.







# AN EASY INTRODUCTION TO CONIC SECTIONS.



## COORDINATES OF A POINT.

1. In Plane Geometry the term *Ordinate* means one of any number of parallel lines drawn from successive points in one plane, to meet a straight line in the same plane perpendicularly or at any angle.

In the following diagrams, OX and OY are intersecting straight lines. The successive lines PL, QM, RN, YO are




drawn *ordinately*, or in parallel succession, to meet OX; and the lines DR, CQ, BP, OX are drawn *ordinately* to OY. In one of the diagrams the parallels are perpendicular to the line they meet; in the other they are oblique.

2. The lines PL, PB, being both drawn from P, are

together called *Coordinates* of that point; in like manner,  $QM$ ,  $QC$  are coordinates of the point  $Q$ ; and so forth.

But the term *ordinate* is usually restricted to the lines that are parallel to  $OY$ ; and instead of those parallel to  $OX$  we use the equivalent magnitudes  $OL$ ,  $OM$ ,  $ON$ , which, as being portions of  $OX$  cut off by the ordinates, are called *abscissæ*, and each of them an *abscissa*. Thus,  $QM$  is the ordinate and  $OM$  the abscissa of the point  $Q$ ; and  $OM$ ,  $QM$  are the coordinates of that point.

3. If, in one plane, two unlimited straight lines,  $X'X$ ,  $YY'$ , be drawn intersecting, at any angle, at the fixed point  $O$ , the position of any other point in the plane may be referred to these lines, and expressed in terms supplied by them. For a point must be on the right or on the left of  $YY'$ , or else in the line  $YY'$ ; and must also be above or below  $X'X$ , or else in the line  $X'X$ . Thus, the point  $P$  is situated on the right of  $Y'Y$ , and above  $X'X$ ; that is, it is within the angle  $YOX$ ; and if the lengths of  $PN$  and  $PM$  be known to be, respectively, 4 and 3, then, measuring  $OM=4$  from  $O$  towards  $X$ , and  $ON=3$  from  $O$  towards  $Y$ , we shall have indicated, by means of  $X'X$  and  $YY'$ , the position of  $P$ , viz. that  $P$ 's position is at the intersection of lines drawn respectively parallel to  $OY$  and  $OX$  from the points  $M$  and  $N$ .



4. The lines  $YOY'$  and  $X'OX$  are called the *Axes of Coordinates*; the former, by itself, the *axis of ordinates*, as being the line on which ordinates are measured; the latter, by itself, the *axis of abscissæ*, as being the line on which abscissæ are measured.


But, in general reasonings, an unknown or variable abscissa is denoted by the letter  $x$ , and its corresponding ordinate by the letter  $y$ ; and therefore it is usual to call the axis of abscissæ the axis of  $x$ , and that of ordinates the axis of  $y$ .

The point O, where the coordinates intersect, is called the *Origin of Coordinates*, because we are to suppose the lines OX, OX', OY, OY' all to be drawn from O; OX to the right, OX' to the left, OY upwards, OY' downwards.

The coordinates are called *Rectangular* or *Oblique*, according as they do or do not intersect at right angles.

In the following part of this introductory treatise, we shall always suppose the axes to be *rectangular*.

5. The position, then, of P is known, if OM and PM are known. Suppose that the lengths of these coordinates are, respectively,  $a$  and  $b$ ; we characterise the position of P thus,  $x=a, y=b$ , which are called equations to the point P.



For shortness, we often speak of the point  $(a, b)$ , meaning the point which has  $a$  for the length of its abscissa, and  $b$  for that of its ordinate. So, the point  $(4, 3)$  means the point whose coordinates are  $x=4, y=3$ .

6. Suppose it is required to determine the geometrical position of the point  $(-10, 7)$ .

Here, since  $x=-10$ , and  $y=7$ , we take  $ON=7$  upward from the origin, and  $OL=10$  to the left of the origin; and, completing the parallelogram OR, we obtain R, the required position, which is within the angle YOX'. This procedure will be readily understood by the student who has read Chap. I. of the Author's *Treatise on Trigonometry*. The point O being the fixed origin of coordinates, an abscissa measured to the right of O, viz. along OX, is accounted positive; one measured to the left of O, viz. along OX', is therefore negative; and, in like manner, an ordinate measured upwards from O, along OY, being accounted positive, one measured downwards from O, viz. along OY', is to be accounted negative.


Suppose the absolute magnitude of  $OL$  = that of  $OM$  = 10, and the absolute magnitude of  $OK$  = that of  $ON$  = 7; then, according to the above conventional understanding, the point  $P$  is (10, 7),  $Q$  is (10, -7),  $R$  is (-10, 7),  $S$  is (-10, -7).

If a point whose abscissa is  $=a$  be situated on the line  $OX$ , then the equations of the point are  $x=a$ ,  $y=0$ , or the point is  $(a, 0)$ . And if a point whose ordinate is  $=b$  be situated on  $OY$ , the equations are  $x=0$ ,  $y=b$ , or the point is  $(0, b)$ . If the point be situated on both axes, it is, of course, their point of intersection, viz. the origin, which is characterised by the equations  $x=0$ ,  $y=0$ .

#### EQUATION TO THE STRAIGHT LINE.

7. If  $OM=a$ , and  $PM=b$ , the coordinates of every point in the unlimited line  $OP$  are in the ratio of  $a : b$ . Thus, by similar triangles,

$$\frac{PM}{OM} = \frac{QL}{OL} = \frac{RN}{ON} = \frac{b}{a}, \text{ or generally } \frac{y}{x} = \frac{b}{a}, \text{ or } y = \frac{b}{a}x;$$



where it should be observed that  $\frac{b}{a}$  is an expression for the tangent of the angle which  $OP$  makes with the axis of  $x$  in the positive direction, that is, of the angular distance of  $OP$  from  $OX$  measured towards  $OY$ .

For the prolongation of OP downwards, if OM' be taken  $= -a$ , then SM'  $= -b$ , and the tangent of S'OX'  $= \frac{-b}{-a} = \frac{b}{a} = \tan \text{POX}$ .

It appears, then, that the equation  $y = \frac{b}{a}x$  is satisfied by every point on the unlimited line S'OR.

If OM'  $= -a$ , and SM'  $= b$ , the coordinates of every point on the unlimited line OS are, as regards absolute magnitude, in the ratio of  $a : b$ ; but if the expression for the ratio is to include reference to the line's position, we must now write  $\frac{y}{x} = \frac{b}{-a}$  or  $-\frac{b}{a}$ , that is,  $y = -\frac{b}{a}x$ ; which signifies that the tangent of the angle which SO makes with the axis of  $x$ , in the *positive direction* (viz. of the angle SOX, not SOX'), is  $= -\frac{b}{a}$ .

For the prolongation of SO downwards, we have OM  $= a$ , and P'M  $= -b$  and the tangent of P'OX  $= \frac{-b}{a}$  or  $-\frac{b}{a} = \tan \text{SOX}$ .

It appears, then, that the equation  $y = -\frac{b}{a}x$  is satisfied by every point on the unlimited line P'OS.

**3.** *To find the general Equation to a Straight Line, referred to rectangular axes.*

The position of a straight line in relation to the axes is determined, when we know the inclination to OX of that part of the line which is above X'X, and also the distance of the line's intersection with YY' from the origin O.

In the last article we found the *particular* equation characterising a straight line that intersects with YY' at the origin. That equation may be written  $y = mx$ ; where  $m$  denotes the tangent of the line's inclination to OX in the *positive direction*.

But, now, let  $AT$  be a straight line, of indefinite length, meeting the axis of  $y$  at  $B$ .

Draw  $OF$ , through the origin, parallel to  $AT$ . In  $x'$   $AT$  take any point  $Q$ ; and draw  $QM$  parallel to  $OY$ , meeting  $OX$  at  $M$ , and  $OF$  at  $P$ .

$$\frac{PM}{OM} = \tan FOX; \therefore PM = OM \tan FOX;$$

$$QM = PM + PQ = PM + OB;$$

$$\therefore QM = OM \tan FOX + OB.$$




Hence, if  $OB$ , which is called the *intercept* on the axis of  $y$ , be given in length  $=c$ , and  $\tan FOX$ , or  $\tan TAX$ ,  $=m$ , and if the coordinates of the point  $Q$  be  $x, y$ , then the equation to the straight line  $AT$  will be

$$y = mx + c.$$

Suppose the line  $AT$  still to cut the axis of  $y$  above  $X'X$ , but to make the angle  $TAX$  obtuse; then the tangent of that angle will be negative, and  $m$  in the above equation stands for a negative quantity,  $c$  being still positive.


Suppose the line  $AT$  to cut the axis of  $y$  below  $X'X$ , and to make the angle  $TAX$  acute; then we have  $QM = PM - PQ = PM - OB$ , or,  $QM = OM \tan FOX - OB$ ; and in this instance the symbol  $m$  stands for a positive quantity, and  $c$  for a negative.

Lastly, suppose the line  $AT$  to cut the axis of  $y$  below  $X'X$ ,



and to make the angle TAX obtuse; we shall then have  $m$  and  $c$  both standing for negative quantities.

Thus, then, if the absolute magnitude of  $m$ , in each of the particular cases supposed, were  $=\frac{b}{a}$ , and that of  $c$  were  $=n$ ,



the four varieties of lines would be represented as follows:

$$(i.) y = \frac{b}{a}x + n. \quad (ii.) y = -\frac{b}{a}x + n.$$

$$(iii.) y = \frac{b}{a}x - n. \quad (iv.) y = -\frac{b}{a}x - n.$$

So long as the same straight line, unrestricted as to length, is considered, the quantities  $m$  and  $c$  represent definite magnitudes, and are therefore called *constants*; while  $x$ , or  $y$ , is arbitrary, and may have an indefinite variety of values, to which  $y$ , or  $x$ , will have the same variety of *corresponding* values; the symbols  $x$  and  $y$ , therefore, are called *variables*.

9. Observe how the general equation

$$y = mx + c$$

provides for the following cases:

If the line pass through the origin of coordinates, then  $c=0$ , and the equation becomes

$$y = m x, \text{ as formerly found.}$$

If the line be parallel to the axis of  $x$ , it makes *no geometrical angle* with that axis; then  $m=0$ , and the equation becomes

$$y = c;$$

which signifies that the ordinate of *any* point on the line is  $=c$ .

If the line be parallel to the axis of  $y$ , then, when the



axes are rectangular,  $m$  is the tangent of a right angle, and is therefore infinite, and the required line will not cut off any intercept from the axis of  $y$ , so that  $c$  also is infinite, and if  $a$  be any abscissa, positive or negative, then  $\frac{c}{a}$ , which is also infinite, is equivalent to  $m$ ; we shall thus have

$$y = \frac{c}{a}x + c, \text{ or } \frac{y}{\frac{c}{a}} = x + a, \text{ or } 0 = x + a,$$

hence, as  $a$  may be positive or negative, the equation will be

$$x = \pm a;$$

which signifies that the abscissa of any point on the line is  $= a$ .

The case, however, of a line parallel to the axis of  $y$  is very simply investigated by itself. For we need only say, Let  $a$  be the abscissa for *one* point on the required line, and  $x$  the abscissa of *any* point on the line; and we have  $x = a$ .

**10.** *The Equation to a Straight Line that makes intercepts on the axes may be expressed in terms of the intercepts.*

Let  $AB$  be a straight line cutting the axes at  $A$  and  $B$ ; and let the intercepts  $OA$ ,  $OB$  be respectively equal to  $a$  and  $b$ ; then  $m$ , the tangent of  $BAX$ ,  $= -\frac{b}{a}$ , and  $c = OB = b$ ; and the equation


$$y = mx + c$$

becomes  $y = -\frac{b}{a}x + b$ , or  $\frac{b}{a}x + y = b$ ;

then dividing by  $b$  gives

$$\frac{x}{a} + \frac{y}{b} = 1, \text{ the required equation.}$$

**11.** It has now become evident that any straight line can be represented by an equation of the first degree.



We can prove also that any equation of the first degree will serve to represent a straight line.

For, an equation of the first degree being one into whose simple form there does not enter any power of an unknown quantity higher than the first power, or any product of unknown quantities, the general form of the indeterminate equation of the first degree is

$$A x + B y + C = 0;$$

where A, B, C may be finite or zero.

When A is zero, we have  $B y = -C$ , or  $y = -\frac{C}{B}$ ; which, as we have seen, represents a straight line parallel to the axis of  $x$ .

When B is zero, we have  $A x = -C$ , or  $x = -\frac{C}{A}$ ; which, as we have seen, represents a straight line parallel to the axis of  $y$ .

If B is not zero, then, dividing by B, we have

$$\frac{A}{B}x + y + \frac{C}{B} = 0, \quad \text{or, } y = -\frac{A}{B}x - \frac{C}{B}.$$

which, as we have seen, represents a line meeting the axis of  $y$  at a distance  $= -\frac{C}{B}$  from the origin, and making with OX an angle whose tangent is  $= -\frac{A}{B}$ .

Therefore the equation  $A x + B y + C = 0$  will always serve to represent a straight line.

**12.** The following statements should now be distinctly understood :

The equation to a line is one that expresses an invariable relation which is satisfied by the coordinates of every point on the line.

The word *locus* (plural *loci*) is used to signify a lineal common-place or path, the coordinates of every point of which have the same ratio.

A line, whether straight or curved, is called the *locus* of a given equation, when the coordinates of every point on the line satisfy the equation.

**13.** It will now be a useful exercise for the student to draw the straight lines represented by a few given equations.

*Example* (1). It is required to draw the straight line whose position is represented by  $2x - 5y = 7$ .

*1st Method.* Make the given equation assume the form


$$y = mx + c;$$

which is done by dividing by 5 and transposing; thus,

$$\frac{2}{5}x - y = \frac{7}{5}; \therefore y = \frac{2}{5}x + (-\frac{7}{5}).$$

Here we have  $m = \frac{2}{5}$  = the value of the tangent of the angle which that part of the required line which is above the axis of  $x$  will make with the axis of  $x$  produced in the positive direction.

First, then, draw the rectangular axes  $X'X, Y'Y$ ; make  $PM$  perpendicular to  $OM$ , so that  $OM : PM$  as 5 : 2; and join  $OP$ .



The angle  $POX$  is that of the required line's inclination to

$OX$ , because the tangent is  $= \frac{PM}{OM} = \frac{2}{5}$ . Now, any line pa-

rallel to  $OP$  will have the same inclination to  $OX$ . But the required line is to make on the axis of  $y$  a *negative* intercept whose absolute length is  $\frac{7}{5}$ . Accordingly, measuring  $OB = \frac{7}{5}$  downwards from  $O$ , and drawing through  $B$  a line parallel to  $OP$ , we have  $AB$  the required line, making with  $OX$  the positive angle  $TAX$ , whose tangent is  $= \frac{QM}{AM} = \frac{PM}{OM} = \frac{2}{5}$ .

*2nd Method.* Make the given equation assume the form

$$\frac{x}{a} + \frac{y}{b} = 1;$$

which is done by dividing by 7, and then putting the coefficients of  $x$  and  $y$  in the form of divisors; thus,

$2x-5y=7$  will become  $\frac{2}{5}x-\frac{1}{5}y=1$ ;

$$\text{or, } \frac{x}{\frac{5}{2}}-\frac{y}{\frac{5}{1}}=1; \text{ or, } \frac{x}{\frac{5}{2}}+\frac{y}{-\frac{5}{1}}=1;$$

where  $a=\frac{5}{2}$ ,  $b=-\frac{5}{1}$ .

If we take, therefore, from OX a positive abscissa, OA, whose absolute length is  $\frac{5}{2}$ , and from OY' a negative ordinate, OB, whose absolute length is  $\frac{5}{1}$ , we shall have two points A and B on the required line, which are, of course, sufficient to determine the position of the line.

*3rd Method.* First, suppose  $y=0$ , and then find the corresponding value of  $x$ , to be used as an abscissa; secondly, suppose  $x=0$ , and find the corresponding value of  $y$ , to be used as an ordinate.

For, since the equation  $2x-5y=7$  represents a line of indefinite length, one point of the line will meet the axis of  $x$ , and then,  $y$  being  $=0$ , we have

$$2x-0=7; \text{ or, } x=\frac{7}{2};$$

also, another point of the line will meet the axis of  $y$ , and then,  $x$  being  $=0$ , we have


$$0-5y=7; \text{ or, } y=-\frac{7}{5}.$$

*Note.* The 2nd and 3rd methods are inapplicable in such a case as  $x-2y=0$ .

*Ex. (2).* Draw the straight line  $y-x\sqrt{2}=5$ .

Here, suppose  $x=0$ , then  $y=5$ ;  
therefore take the positive ordinate  
OB=5.

Again, suppose  $y=0$ , then  $x=-5+\sqrt{2}=-\frac{5}{2}\sqrt{2}$ ; therefore take the negative abscissa OA= $\frac{1}{2}(5\sqrt{2})$ , which is conveniently done thus: make OM=OB=5; draw MD perpendicular and equal to OM, and produce it to P, so that PM may be equal to the hypotenuse OD= $5\sqrt{2}$ ; then OA= $\frac{1}{2}$ PM. Hence, to draw the required line, join AB.



## EXERCISES [A].

Draw the lines represented by the following equations :

- |   |                                       |
|---|---------------------------------------|
| 1. $3x - 2y = 4.$                               | 2. $x + 3 = 4y.$                      |
| 3. $4x + 5y = 6.$                               | 4. $\frac{1}{2}y - \frac{1}{3}x = 5.$ |
| 5. $3x + 4y + 5 = 0.$                           | 6. $x + y = 0.$                       |
| 7. $x\sqrt{3} - y = 8.$                         | 8. $x = -2.$                          |
| 9. $\frac{4}{3}x + \frac{2}{3} = \frac{1}{2}y.$ | 10. $6y + x\sqrt{5} + \sqrt{10} = 0.$ |

## PROBLEMS ON THE STRAIGHT LINE.

**14.** *To find the Equation to a Straight Line that shall pass through a given point  $(x', y')$ \*.*

Taking  $x$  and  $y$  as the coordinates of an indeterminate point on the line, let the line be represented by

$$y = mx + c. \quad (1.)$$

The given coordinates  $x', y'$  must satisfy the general equation ; therefore,

$$y' = mx' + c. \quad (2.)$$

hence, eliminating  $c$  by subtraction, we have

$$y - y' = m(x - x'), \quad (3.)$$

which is the required equation, and which may be written, if we choose, in the form of the general equation, thus :

$$y = mx + y' - mx';$$

where  $m$  is perfectly arbitrary.

It is evident that the required equation must reject either  $m$  or  $c$ ; for it is in general impossible to draw a straight line through a given point so as to meet the axis of  $y$  at a proposed distance  $c$  from the origin, and at the same time make with the axis of  $x$  a proposed angle,  $\tan^{-1}m$ .

If we choose,  $m$  may be rejected and  $c$  retained. Thus, from equation (2) we have  $m = \frac{y' - c}{x'}$ , which will convert (1) into the required form

$$y = \frac{y' - c}{x'}x + c.$$

\* Given coordinates of a point are very often expressed by the letters  $x$  and  $y$  accented.

**15.** To find the equation to a straight line that shall pass through two given points,  $(x', y')$  and  $(x'', y'')$ .

Here we can assume the equations (1), (2), (3) from the preceding article :

$$y = mx + c, \quad (1.)$$

$$y' = mx' + c, \quad (2.)$$

$$y - y' = m(x - x') \quad (3.)$$

(3) representing a line that fulfils one of the conditions now proposed : that is, a line passing through the point  $(x', y')$ .

The line, however, is also to pass through  $(x'', y'')$ , and, therefore,

$$y'' = mx'' + c. \quad (4.)$$

Subtracting (2) from (4) we have

$$y'' - y' = m(x'' - x'),$$

$$\text{whence, } m = \frac{y'' - y'}{x'' - x'};$$

$m$  being thus no longer arbitrary.

Substituting in (3) the value of  $m$  just found :

$$y - y' = \frac{y'' - y'}{x'' - x'}(x - x') \quad (5.)$$

is the required equation. It may be written, if we choose, in the form of the general equation, thus :

$$y = \frac{y'' - y'}{x'' - x'}x + \frac{y'x'' - x'y''}{x'' - x'}.$$

If we suppose  $x'' = x'$ , the line will be parallel to the axis of  $y$ ; for the tangent will be  $= \frac{y'' - y'}{0} = \infty$ , = the tangent of a right angle.

If we suppose  $y'' = y'$ , the line will be parallel to the axis of  $x$ ; for the tangent becomes  $= \frac{0}{x'' - x'} = 0$ , = the tangent of the trigonometrical angle which  $O X'$  makes with  $O X$ .

If we suppose  $x' = 0$ ,  $y' = 0$ , we make the point  $(x', y')$  the origin; for which case the equation is reduced to the simple form  $y = \frac{y''}{x''}x$ .

**16.** To find the equation to a straight line which shall pass through a given point, and be parallel to a given straight line.

Let the given line be represented by the equation

$$y = m'x + c', \quad (1.)$$

where  $m'$  and  $c'$  are known;

and let the equation to the required line be

$$y = mx + c, \quad (2.)$$

where  $m$  and  $c$  are not known.

Then, since the lines represented by these equations are parallel, they must have the same inclination to OX, that is,  $m = m'$ ; thus the second equation becomes

$$y = m'x + c, \quad (3.)$$

where  $c$ , the intercept on the axis of  $y$ , remains indeterminate, because an infinite number of lines may be parallel to a given line. The equation, in effect, asserts that it matters not what the intercept on the axis of  $y$  may be, provided the tangent of the inclination of the given line to the axis of  $x$  be also the tangent for any other line we may draw.

But, if the parallel be restricted to pass through a given point  $(x', y')$ , then

$$y' = m'x' + c; \quad (4.)$$

Hence, subtracting (4) from (3), we have

$$y - y' = m' (x - x') \quad (5.)$$

which is the required equation.

**17.** To find an expression for the tangent of  $\phi$ , the angle of intersection of two given straight lines.

Let AA' and CC' be the two given lines, intersecting at P.

Let the equation to AA' be


$$y = m'x + c'$$

and the equation to CC'

$$y = m''x + c'';$$

$m'$  is the tangent of the angle PAX;

$m''$  is the tangent of the angle PCX.



Now,  $\tan APC = \tan (PCX - PAX)$

$$= \frac{\tan PCX - \tan PAX}{1 + \tan PCX \tan PAX} = \frac{m'' - m'}{1 + m'm''};$$

$$\therefore \tan \text{ of supplement } A'PC = \frac{m' - m''}{1 + m'm''},$$

Hence, the required expression is  $\tan \phi = \pm \frac{m'' - m'}{1 + m'm''}$

according as the angle is to the left or the right of  $CC'$ .

When the lines become parallel, the tangent = 0, and therefore the numerator in the above expression = 0, that is  $m'' = m'$ .

In order that the angle of intersection may be a right angle, the tangent =  $\infty$ , and therefore the denominator in the above expression = 0, or  $m'm'' = -1$ , or  $m'' = -\frac{1}{m'}$ .

Hence, 
$$y = -\frac{1}{m'}x + c''$$

represents a line perpendicular to a given line whose position is expressed by  $y = m'x + c'$ .

Of course, there are an infinite number of such perpendiculars. Two lines intersect at right angles whenever the coefficient denoting the tangent in the equation to one line is the reciprocal, with a contrary sign, of that in the equation to the other line. In the next problem we fix one of the lines by the condition of its passing through a given point.

**18.** *To find the equation to a straight line that shall pass through a given point, and be perpendicular to a given straight line.*

Let the coordinates of the given point be  $x', y'$ , and let the equation to the given line be

$$y = mx + c.$$

The equation to a line through  $(x', y')$  is of the form

$$y - y' = m'(x - x');$$



and if this equation is perpendicular to the former, then  $m' = -\frac{1}{m}$ , and the required equation is

$$y - y' = -\frac{1}{m}(x - x').$$

**19.** To find the equation to a straight line that passes through a given point  $(x', y')$ , and makes a given angle  $\phi$  with a given straight line.

Let the equation to the given line be

$$y = mx + c; \quad (1.)$$

the equation to the required line passing through  $(x', y')$  will be of the form

$$y - y' = m'(x - x'). \quad (2.)$$

By art. 17, we have  $\tan \phi = \pm \frac{m - m'}{1 + mm'}$ ;

hence,  $m' = \frac{m - \tan \phi}{1 + m \tan \phi}$  or  $\frac{m + \tan \phi}{1 - m \tan \phi}$ ;

therefore (2) becomes

$$y - y' = \frac{m - \tan \phi}{1 + m \tan \phi} (x - x'). \quad (3.)$$

or else,  $y - y' = \frac{m + \tan \phi}{1 - m \tan \phi} (x - x'). \quad (4.)$

Either of the equations (3) and (4) will be the one required; for generally through a given point two straight lines can be drawn, making a given angle with a given straight line.

*Note* (1). When  $m = 0$ , the given line is parallel to the axis of  $x$ ; then the required equations are

$$y - y' = \pm \tan \phi (x - x').$$

*Note* (2). When  $m = \infty$ , the given line is parallel to the axis of  $y$ . For this case, dividing numerator and denominator of  $\frac{m + \tan \phi}{1 - m \tan \phi}$  by  $m$ , we have

$$\frac{1 + \frac{\tan \phi}{m}}{\frac{1}{m} - \tan \phi} = \frac{1 + \frac{\tan \phi}{\infty}}{\frac{1}{\infty} - \tan \phi} = \frac{1 + 0}{-\tan \phi} = -\frac{1}{\tan \phi}$$

Similarly,  $\frac{m - \tan \phi}{1 + m \tan \phi}$  becomes  $= \frac{1}{\tan \phi}$ .

Then the required equations are

$$y - y' = + \cot. \phi (x - x').$$

**Note (3).** Dividing numerator and denominator by  $\tan \phi$ ,

we have  $\frac{\frac{m}{\tan \phi} + 1}{\frac{1}{\tan \phi} + m}$ , which, when  $\phi$  is a right angle, is

$$\frac{\frac{m}{\infty} \pm 1}{\frac{1}{\infty} \mp m} = \frac{0 \pm 1}{0 \mp m} = -\frac{1}{m},$$


as found in art 18.

Or thus,  $\frac{\frac{m}{\tan \phi} + 1}{\frac{1}{\tan \phi} + m} = \frac{m \cot \phi + 1}{\cot \phi + m} = -\frac{1}{m};$

the cotangent of a right angle being zero.

**20.** To express the distance between two points in terms of their coordinates.

Let P and Q be two points whose coordinates are given. Let the coordinates of Q be  $x', y'$ , and let those of P be  $x'', y''$ . Join PQ; from P and Q draw PM, QN parallel to OY, and draw QR parallel to OX.



We have  $\angle PRQ$  a right angle ; therefore (by Euclid, I. 47)

$$PQ^2 = QR^2 + PR^2.$$

Now,  $QR = OM - ON = x'' - x'$ ; and  $PR = PM - QN = y'' - y'$ ; hence  $PQ^2 = (x'' - x')^2 + (y'' - y')^2$ ; which determines  $PQ$ .

And here it should be observed, that, as only the absolute length of PQ is required, we need take no notice of the double sign  $\pm$  that would distinguish the algebraic roots of the value of PQ<sup>2</sup>.


If the point Q were at the origin, we should have  $x'=0$ ,  $y'=0$ ; and then

$$PQ^2 = (x''-0)^2 + (y''-0)^2 = x''^2 + y''^2.$$

And, if Q were on the axis of  $x$ , and P on that of  $y$ , we should have  $x'=0$ ,  $y'=0$ ; and then

$$PQ^2 = (0-x')^2 + (y''-0)^2 = x'^2 + y''^2.$$

Again, let P and Q be two points; let the given coordinates of Q be  $-a$ ,  $b$ , and let those of P be  $h$ ,  $k$ . Then, the expression for  $PQ^2$ , corresponding to that found above, will be  $PQ^2 = \{h - (-a)\}^2 + (k - b)^2$ ,  
 $= (h+a)^2 + (k-b)^2$



**21.** To determine the distance of a given point from a given straight line.

Let P be the given point, and AB the given line; it is required to determine the length of the perpendicular PR.

If the coordinates of P be  $h$ ,  $k$ , and those of R be  $x_1$ ,  $y_1$ , the analytical expression for the length of PR is

$$PR^2 = (x_1 - h)^2 + (y_1 - k)^2; \quad (1.)$$

Let the equation to the given line ABR be

$$y = mx + c;$$

then the equation to PR, perpendicular to ABR, is

$$y - k = -\frac{1}{m}(x - h);$$

and, as R is a point common to both these lines, we have


$$y_1 - mx_1 + c, \quad (2.)$$

$$y_1 - k = -\frac{1}{m}(x_1 - h). \quad (3.)$$

Multiplying (3) by  $m^2$ , and then adding (2), we have

$$(1 + m^2)y_1 - m^2k = -mh + c;$$

$$\text{whence, } y_1 = \frac{m^2k + mh + c}{1 + m^2}.$$



Multiplying (2) and (3) by  $m$ , and then subtracting, we get

$$mk = (1 + m^2)x_1 - h + mc;$$

$$\text{whence, } x_1 = \frac{mk + h - mc}{1 + m^2}.$$

Hence, by substitution in (1),

$$\begin{aligned} PR^2 &= \left( \frac{mk + h - mc}{1 + m^2} - h \right)^2 + \left( \frac{m^2k + mh + c}{1 + m^2} - k \right)^2 \\ &= \left( \frac{mk + h - mc - h - m^2h}{1 + m^2} \right)^2 + \left( \frac{m^2k + mh + c - k - m^2k}{1 + m^2} \right)^2 \\ &= \frac{m^2}{(1 + m^2)^2} (k - mh - c)^2 + \frac{1}{(1 + m^2)^2} (k - mh - c)^2 \\ &= \frac{1}{1 + m^2} (k - mh - c)^2; \end{aligned}$$

$$\therefore PR = \frac{k - mh - c}{\sqrt{1 + m^2}}, \text{ the expression required.}$$

The positive and negative values of the denominator refer to the ambiguity of the direction of PR, as P may be on either side of the given line; but, as only the magnitude of PR is required, the denominator may be taken with the sign that will give a positive value for PR.

**22.** We shall now subjoin a few Examples and Exercises on articles 14-21.


### EXAMPLES.

*Ex. (1).* Draw, and represent by an equation, the straight line passing through the points P, Q, which are  $(-4, 5)$  and  $(7, -1\frac{1}{2})$ , respectively. Find also the length of PQ.

Take OM, PM, = 4, 5; ON, QN, = 7,  $1\frac{1}{2}$ ; the line PQ, indefinitely produced, is the line required.

From art. 15, we have

$$y - 5 = \frac{-1\frac{1}{2} - 5}{7 + 4}(x + 4),$$



$$\text{or, } y-5 = -\frac{1}{2}(x+4),$$

which gives  $13x+22y=58$ , the required equation.

The equation determining the length PQ (*see* art. 20)

$$\begin{aligned} \text{is } PQ^2 &= (-4-7)^2 + (5+1\frac{1}{2})^2 \\ &= 121 + 1\frac{1}{4} \cdot 9 = 6\frac{1}{4} \cdot 3; \therefore PQ = \frac{1}{2}\sqrt{653}. \end{aligned}$$

*Ex. (2).* What will be the equation to a straight line passing through the point (21, 7) at right angles to the line  $y-3x=24$ ?

(*See* art. 18.) We have here  $x'=21$ ,  $y'=7$ ; and since the given line is represented by  $y=3x+24$ , we have  $m=3$ .

Hence,  $y-7 = -\frac{1}{3}(x-21)$ , or  $3y-21 = -x+21$ ,  
gives  $x+3y=42$ , the required equation.

*Ex. (3).* Shew that the straight lines

$$4x+3y=6, \quad 5x+4y=7, \quad 2x-y=8,$$

intersect at one point.

Here, by means of any two of the given equations, we can find values of  $x$  and  $y$  satisfying both. Take the 1st and 2nd:

$$\begin{array}{r} 5x=7-4y \\ 4x=6-3y \\ \hline \end{array}$$

By subtraction,  $x=1-y$

$$\therefore 5(1-y)=7-4y, \text{ or } y=-2, \text{ and } 1-y=x=3.$$

Therefore, the coordinates of the point of intersection of the 1st and 2nd lines are  $x=3$ ,  $y=-2$ ; and these will be found to be coordinates of a point on the 3rd line also; for  $2x-y=6+2=8$ .

*Ex. (4).* If the line  $y=mx+7$  pass through the intersection of the lines  $y=2x+6$  and  $3y=x-7$ , what is the magnitude of  $m$ ?

$$3y=6x+18=x-7, \text{ or } x=-5,$$

$$2x+6=y=-4;$$

$$\text{hence, } mx+7=-4,$$

$$\text{that is, } -5m+7=-4; \text{ or, } m=\frac{1}{5}.$$

*Ex. (5).* Find the distance of the origin of rectangular coordinates from the line  $\frac{x}{a} + \frac{y}{b} = 1$ .

The given equation may be put in the form

$$y = -\frac{b}{a}x + b;$$


where we have  $m = -\frac{b}{a}$ ; hence, applying the result found

in art. 21, we have  $h=0$ ,  $k=0$ ,  $c=b$ ; and therefore

$$\begin{aligned} \text{OR} &= \frac{k-mh-c}{\sqrt{(1+m^2)}} = -b \div \sqrt{\left(1 + \frac{b^2}{a^2}\right)} \\ &= -ab \div \sqrt{(a^2 + b^2)}; \end{aligned}$$

or the magnitude of the required distance is

$$= \frac{ab}{\sqrt{(a^2 + b^2)}}.$$



*Note.* Since the triangle  $OAB = \frac{1}{2}$  the rectangle  $OA \cdot OB$ , and also  $= \frac{1}{2}$  the rectangle  $AB \cdot OR$ , we have  $AB \cdot OR = OA \cdot OB$ ;

but  $OA=a$ ,  $OB=b$ , and  $AB = \sqrt{(a^2 + b^2)}$ ;


hence,  $OR \sqrt{(a^2 + b^2)} = ab$ ; and  $OR = ab \div \sqrt{(a^2 + b^2)}$ ; which verifies the formula.

*Ex. (6).* What points on the axis of  $x$  are at a perpendicular distance  $a$  from the line

$$\frac{x}{a} + \frac{y}{b} = 1?$$

Take  $OA=a$ ,  $OB=b$ ; then  $AB$  is the given line.

Let  $PD$ ,  $P'E$  be the perpendicular distances of  $AB$  from  $OX$ , and each  $=a$ .



$AB = \sqrt{(a^2 + b^2)}$ ; also, by similar triangles,  $\frac{AP}{PD} = \frac{AB}{OB}$ ,

$$\text{whence, } AP = \frac{a}{b} \sqrt{(a^2 + b^2)}.$$

Therefore, the abscissa of the point P,  $= OA + AP$ , is

$$a + \frac{a}{b} \sqrt{(a^2 + b^2)};$$

and, as  $OP'$  is on the negative side of  $O$ , therefore the abscissa of the point  $P'$ ,  $=OA - AP$ , is

$$a - \frac{a}{b} \sqrt{a^2 + b^2}.$$


*Ex. (7).* The coordinates of a point  $P$  are 4 and 10, and those of another point  $Q$  are 17 and 5; find the coordinates of a point  $R$  in the straight line  $PQ$ , making  $PR : QR :: 3 : 4$ .

Let  $x, y$  be the coordinates of  $R$ ;  
 $ML : NL :: PR : QR :: 3 : 4$ ;

$$\frac{ML}{NL} = \frac{x-4}{17-x} = \frac{3}{4}; \therefore x = 9\frac{4}{7}.$$


Again,  $PQ : QR :: 7 : 4 :: HP : KR$ ;  $\therefore KR = \frac{4}{7}HP = \frac{4}{7}(10-5) = 2\frac{4}{7}$ ; hence  $y = KR + QN = 2\frac{4}{7} + 5 = 7\frac{6}{7}$ .

Therefore the coordinates of  $R$  are  $9\frac{4}{7}, 7\frac{6}{7}$ .



*Ex. (8).* Find the area of a triangle, the coordinates of its angular points being 8 and 9, 13 and 21, 23 and 12, for  $A, B, C$ , respectively.

Here we have  $OL, AL = 8, 9$ ;  
 $OM, BM = 13, 21$ ;  
 $ON, CN = 23, 12$ .



Area of trapezium  $LABM$

$$= \frac{1}{2}(AL + BM)LM = \frac{1}{2}(9 + 21)(13 - 8) = 75$$

Area of trapezium  $MBCN$

$$= \frac{1}{2}(BM + CN)MN = \frac{1}{2}(21 + 12)(23 - 13) = 165$$

$$\therefore \text{Area of } LABCN = 240$$

$$\text{Area of trapezium } LACN = \frac{1}{2}(9 + 12)(23 - 8) = 157\frac{1}{2}$$

$$\therefore \text{Area of } ABC = 82\frac{1}{2}$$

*Ex. (9).* Given  $BC$ , the base of a triangle,  $= a$ , and the difference of the squares of its sides  $= d^2$ , to determine the locus of the vertex.


Take the base as the axis of  $x$ , and its middle point  $O$  as the origin; and let the coordinates of the vertex  $A$  be  $x$  and  $y$ .

$$BA^2 = BD^2 + DA^2 = (\frac{1}{2}a - x)^2 + y^2,$$

$$AC^2 = DC^2 + DA^2 = (\frac{1}{2}a + x)^2 + y^2;$$

$$\text{hence } AC^2 - BA^2 = 2ax = d^2;$$

$$\therefore x = \frac{d^2}{2a}.$$



This is the equation to a straight line parallel to the axis of  $y$  (see art. 9). Therefore the locus (see art. 12) of the vertex  $A$  is a perpendicular to the base, at the distance  $\frac{d^2}{2a}$  from the middle of the base.

*Ex. (10).* Find the equations to the diagonals of a quadrilateral whose sides are represented by the equations

$$y = x, y = 3x, x = 3, y = 5.$$

Take  $OA = 3$ ,  $AP = 3$ ,  $AQ = 5$ ,  $AB = 9$ ; join  $OP$ ,  $OQ$ ,  $OB$ ; draw  $QR$  parallel to  $OA$ , and join  $PR$ . Then  $OPQR$  is the quadrilateral; for  $PQ$  is the line  $x = 3$ , also  $QR$  is the line  $y = 5$ ,  $OR$  is represented by  $y = 3x$ , and  $OP$  by  $y = x$ .

$$\text{Now, } \frac{OM}{RM} = \frac{OA}{AB} = \frac{1}{3}; \therefore OM = \frac{1}{3}RM = 1\frac{2}{3}.$$

Wherefore, the coordinates of  $R$  are  $1\frac{2}{3}$ ,  $5$ , and those of  $P$  are  $3$ ,  $3$ ; hence the equation to the diagonal  $PR$  is


$$y - 5 = \frac{3 - 5}{3 - 1\frac{2}{3}} (x - 1\frac{2}{3});$$

$$\text{or } y - 5 = -\frac{3}{2}(x - 1\frac{2}{3}), \text{ or } 2y - 10 = -3x + 5,$$

$$\text{which gives } 2y + 3x - 15 = 0.$$

Also, the coordinates of  $O$  are  $0, 0$ , and those of  $Q$  are  $3, 5$ ; hence the equation to the diagonal  $QO$  is simply

$$3y = 5x.$$



*Ex. (11).* Shew that in the figure to Euclid, I. 47, the lines  $BK$ ,  $CF$ ,  $AL$  meet at one point.



Take A as the origin, AB, AC, produced indefinitely, as the axes of  $x$  and  $y$ .

Let  $AB=a$ ,  $AC=b$ .

The equation to BC is  $y = -\frac{b}{a}x + b$ .

The equation to AL, passing through the origin, perpendicular to BC, is  $y = \frac{a}{b}x$ . (1)

The equation to BK, passing through the points  $(-b, b)$  and  $(a, 0)$ , is  $y - b = \frac{0-b}{a+b}(x+b)$ ,

$$\text{or, } y = -\frac{b}{a+b}x + \frac{ab}{a+b}. \quad (2)$$

The equation to CF, passing through the points  $(0, b)$  and  $(a, -a)$ , is  $y - b = \frac{-a-b}{a}(x-0)$ ,

$$\text{or, } y = -\frac{a+b}{a}x + b. \quad (3)$$

We are now to shew that the lines (1), (2), (3) intersect at one point.

The values of  $x$  and  $y$  satisfying (1) and (2) will be the coordinates of the point of intersection of these lines, and can be found thus :

$$y = \frac{a}{b}x = -\frac{b}{a+b}x + \frac{ab}{a+b}; \text{ or, } \left(\frac{a}{b} + \frac{b}{a+b}\right)x = \frac{ab}{a+b};$$


$$\text{or, } \frac{a^2 + ab + b^2}{b}x = ab; \text{ whence } x = \frac{ab^2}{a^2 + ab + b^2}.$$

$$\frac{a}{b}x = y = \frac{a^2b}{a^2 + ab + b^2}.$$

Substituting this value of  $y$  in (3), we shall have, if the three lines intersect at one point,

$$\frac{a^2b}{a^2 + ab + b^2} = -\frac{a+b}{a}x + b; \text{ or, } \frac{-b^2(a+b)}{a^2 + ab + b^2} = -\frac{a+b}{a}x;$$

$$\text{or, } x = \frac{b^2}{a^2 + ab + b^2} + \frac{1}{a} = \frac{ab^2}{a^2 + ab + b^2}, \text{ as before.}$$



## EXERCISES [B].

1. Draw the straight line passing through the points (3, -5) and (-5, 2); and represent it by an equation.

2. Draw, and represent by an equation, the straight line passing through the points (0, -12) and (11, 7).

3. What are the equations to two straight lines, one of which passes through the origin, the other through the point (13, 4), and both of them parallel to another line whose equation is  $7y - 3x = 36$ ?

4. Draw, and represent by an equation, a straight line passing through the point (5, -1), and perpendicular to the line  $3y + 5x = 4$ .

5. What are the equations to the straight lines passing through the point (5, 30), and each inclined at an angle of  $60^\circ$  to the line  $\frac{x}{32} + \frac{y}{14} = 1$ ?

6. The coordinates of a point P are 5 and -4, and those of another point Q are -7 and 12. What is the length of the straight line PQ?

7. Find the distance between the points (-12, 15) and (16, 36).

8. Shew that the three lines

$$y = x + 1, \quad 2y = 3x + 18, \quad 3y = 2x - 13,$$

meet at one point.

9. What number for  $m$ , in the equation  $y = -mx + 8$ , would make that equation represent a line passing through the point of intersection of the lines

$$y = \frac{1}{2}x + 3, \quad y = 6x - 12?$$

10. Find the equation to a straight line which passes through the origin, and also through the point of intersection of the lines

$$6x + 7y + 2 = 0, \quad 7x + 9y - 1 = 0.$$

11. The angular points of the triangle ABC are (-21, -25), (5, 26), and (46, 0). Draw the figure, and find the area of the triangle.

12. ABC is a triangle: suppose the points A, B, C, to be,

respectively, (17, 20), (44, 33), (51, 8) : Find the area of the triangle.

13. OPQ is a triangle : Take the point O as the origin of coordinates ; then

(i.) Express the area of the triangle in terms of the co-ordinates of P and Q.

(ii.) Suppose the points P and Q to be respectively (5, 12) and (12, 9), and find the lengths of the three sides.

14. Find the points of intersection of the straight lines

$$x+3y=11, \quad 2x+y=12, \quad 5x+2y=16;$$

and shew that the area of the triangle formed by joining these points is  $32\frac{1}{2}$ .

15. The coordinates of A, B, C, the angular points of a triangle, are, respectively (3, 4), (8, 9), (10, 6). Find the equation to the line DE joining the middle points of AB and BC. Find also the inclination of DE to AC.

16. Find the equations to the diagonals of the rectangle formed by the four lines  $x=a$ ,  $x=a'$ ,  $y=b$ ,  $y=b'$ .

17. Given  $4y=3x$ , and  $x=12$ , the respective equations to OP, PQ two sides of the parallelogram OPQR; and  $12y=23x$ , the equation to the diagonal OQ : Find the equations to the other sides and diagonal ; and also the area of the parallelogram.

18. ABCD is a square, the rectangular coordinates of the points A and C being (2, 3) and (3, 5) : find the equations to the four sides.

19. Find the equation to a straight line bisecting P, the vertical angle of the triangle OPQ ; the equation to the side

OP being  $5y=12x$ , and that to PQ being  $\frac{x}{24} + \frac{y}{7} = 1$ .

20. In the figure of Euclid, I. 47, if KH and FG be produced to meet at P, and PA produced to meet BC at R : shew that PR is perpendicular to BC.

21. It is required to determine the sides and angles of the triangle ABC, the base AB and a perpendicular to it

through A being assumed as the axes of  $x$  and  $y$ , and the equations to BC and AC being, respectively,


$$y = mx + c, \quad y = m'x.$$

## EQUATION TO THE CIRCLE.

**23.** We are now passing from the consideration of rectilinear to that of curvilinear loci; and begin with the circle, as the simplest of the regular curves.

An equation to the circle is one of which the circumference of a circle is the locus.



Now, the circumference of a circle will evidently be characterised by an expression denoting that every point in the curve is equidistant from a fixed point. If, in the annexed figure,  $OP' = OP'' = OP'''$ , and the accented letters denote successive positions of the point P, then the locus of P is the circumference of a circle; and if O be the origin of coordinates of P, we have the sum of the squares of the coordinates constantly equal to the square of radius. Hence, if  $c$  be the radius of the circle,



$$x^2 + y^2 = c^2$$

is an equation to the circle implying the centre to be the origin of coordinates.

**24.** Let us take for the origin of coordinates any point O in the plane of the circle.



Let C be the centre of the circle: its coordinates  $ON = a$ ,

$NC=b$ ; and let  $P$  be any point on the circumference: its coordinates  $OM=x$ ,  $MP=y$ . Draw  $CQ$  parallel to  $OX$ .

Then  $CQ=OM-ON=x-a$ ;  $PQ=PM-CN=y-b$ .

By Eucl. I. 47,  $CQ^2 + PQ^2 = CP^2$ ;

that is,  $(x-a)^2 + (y-b)^2 = c^2$ ,

or, by development,

$$x^2 + y^2 - 2ax - 2by + a^2 + b^2 - c^2 = 0;$$

which is the general equation to the circle, referred to rectangular axes.

When the origin is a point on the circumference, then

$$ON^2 + NC^2 = OC^2 = c^2, \text{ or } a^2 + b^2 - c^2 = 0;$$

and the general equation gives

$$x^2 + y^2 - 2ax - 2by = 0.$$

If  $OX$  coincide with  $CQ$ , and the origin be on the circumference, the centre  $C$  will be the point  $(a, 0)$ , and  $a$  will be  $=c$ , and  $\therefore a^2 - c^2 = 0$ ; then the general equation gives

$$x^2 + y^2 - 2ax = 0.$$

If  $OY$  coincide with  $NC$ , and the origin be on the circumference, the centre  $C$  will be the point  $(0, b)$ , and  $b$  will be  $=c$ , and  $\therefore b^2 - c^2 = 0$ ; then the general equation gives

$$x^2 + y^2 - 2by = 0.$$

But the centre is most frequently taken as the origin; in which case  $a=0$ ,  $b=0$ ; then the general equation gives


$$x^2 + y^2 = c^2, \text{ as in art. 23.}$$

**25.** It is evident, then, that the general equation to the circle, for rectangular axes, is of the form

$$x^2 + y^2 + Ax + By + C = 0;$$

which is an indeterminate equation of the 2nd degree, having unity as the coefficient of each of the squares of  $x$  and  $y$ , and not containing the product of these variables.

And, conversely, it may be shewn that the locus of an equation of the above form is always the circumference of



a circle; although it will include the peculiarities of a circumference reduced to a mere point, and of a curve merely imaginary.

For, making  $x^2 + Ax$  and  $y^2 + By$  complete squares, we may write the above equation thus :

$$x^2 + Ax + \frac{1}{4}A^2 + y^2 + By + \frac{1}{4}B^2 = \frac{1}{4}A^2 + \frac{1}{4}B^2 - C;$$

that is,  $(x + \frac{1}{2}A)^2 + (y + \frac{1}{2}B)^2 = \frac{1}{4}(A^2 + B^2 - 4C);$

which, if compared with the form

$$(x-a)^2 + (y-b)^2 = c^2,$$

will be observed to represent a circle the coordinates of whose centre are  $-\frac{1}{2}A$  and  $-\frac{1}{2}B$ , and of which the radius is  $\frac{1}{2}\sqrt{(A^2 + B^2 - 4C)}$ .

The circumference will be reduced to a mere point, viz. the centre, if  $A^2 + B^2 = 4C$ , for then the sum of the squares on the left side of the equation will be zero. It is more accurate to say here that the circumference will be reduced to one whose radius is indefinitely small.

The locus of the equation will be impossible if  $4C$  be greater than  $A^2 + B^2$ ; for then  $\sqrt{(A^2 + B^2 - 4C)}$  will be the square root of a negative quantity, which is merely imaginary.

**26.** The preceding article indicates an easy method of constructing any circle that is represented by a given equation. Suppose, for instance, it is required to construct the circle whose equation is

$$x^2 + y^2 + 12x - 10y - 60 = 0.$$

Completing squares with  $x^2 + 12x$  and  $y^2 - 10y$ , we have


$$(x+6)^2 + (y-5)^2 = 60 + 6^2 + 5^2 \\ = 121;$$

and comparing this with the form

$$(x-a)^2 + (y-b)^2 = c^2,$$

we see that  $a = -6$ ,  $b = 5$ ,  $c = 11$ .

Accordingly, the circle will be represented, in magnitude and position, as in the annexed figure, where  $ON$  is taken  $= 6$ ,  $NC = 5$ ,  $CP = 11$ .



## TANGENT AND NORMAL TO A CIRCLE.

**27.** A secant to a circle is a straight line meeting the circumference at two points. Suppose one of the points, at which a secant meets the circumference, to remain fixed, while the secant turns round that point and shifts the second point up to the first, the secant in its limiting position is called the tangent to the circle at the first point. (See figure to art. 28.)

The tangent may thus be regarded as a secant passing through *two coincident points* of the circumference; and this consideration suggests an easy method of investigating the equation that will represent the tangent to a circle.

**28.** To find the equation to the tangent at a given point on the circumference of a circle.

Take the centre as origin, that is, let  $x^2 + y^2 = c^2$  be the equation to the circle; and let  $x', y'$  be the coordinates of the point P, on the circumference, at which the tangent is drawn,  $x'', y''$  the coordinates of another point P' on the circumference, near the given point.

The equation to the straight line PP' is

$$y - y' = \frac{y'' - y'}{x'' - x'}(x - x'). \quad (1.)$$

Now,  $x'^2 + y'^2 = c^2$ ; also,  $x''^2 + y''^2 = c^2$ ;

$$\therefore y'^2 - y''^2 = x'^2 - x''^2,$$


that is,  $(y' + y'')(y' - y'') = (x' + x'')(x' - x'')$ ;

$$\therefore \frac{y' - y''}{x' - x''} = \frac{x' + x''}{y' + y''}; \text{ or, } \frac{y'' - y'}{x'' - x'} = -\frac{x' + x''}{y' + y''};$$

hence, by substitution in the equation to PP',

$$y - y' = -\frac{x' + x''}{y' + y''}(x - x'). \quad (2.)$$

Now, when the point P', by the revolution of the straight line P'P round P, has shifted its position on the curve until



the chord  $PP'$  becomes indefinitely small, that is, when the point  $(x'', y'')$  coincides with the point  $(x', y')$ , we have  $x'' - x' = 0$ , and  $y'' - y' = 0$ ; or  $x'' = x'$ , and  $y'' = y'$ ; in which case equation (2) becomes

$$y - y' = -\frac{2x'}{2y'}(x - x');$$

or, the equation to the tangent at the point  $P$ , that is, at the point  $(x', y')$ , is

$$y - y' = -\frac{x'}{y'}(x - x'). \quad (3.)$$

It will now be seen that for the case in which  $x'' = x'$  and  $y'' = y'$ , equation (1) would have given an indeterminate expression, because that equation does not include the condition of  $(x', y')$  and  $(x'', y'')$  being points on the circumference of a circle.

The equation to the radius  $CP$  is, of course,

$$y = \frac{y'}{x'}x;$$

compare it with (3): the tangent number in one is the reciprocal, with contrary sign, of that in the other, thus shewing that a tangent to a circle is always perpendicular to the radius passing through the point of contact.

Equation (3) may be changed to a simpler form; for by reduction we have

$$\begin{aligned} yy' - y'^2 &= -xx' + x'^2, \\ \text{or, } xx' + yy' &= x'^2 + y'^2, \\ \text{that is, } xx' + yy' &= c^2. \end{aligned}$$

This is the form in which the equation to the tangent is most frequently referred to.

When the equation to the circle is in the form

$$(x - a)^2 + (y - b)^2 = c^2,$$

the equation to the tangent at any point  $(x', y')$  may be similarly found to be

$$(x - a)(x' - a) + (y - b)(y' - b) = c^2.$$

**29.** To express the tangent to a circle in terms of the tangent of the angle which the line makes with the axis of  $x$ .

We have seen that the tangent of the angle which  $QP$ ,



the tangent to the circle, makes with the axis of  $x$  is  

$$= -\frac{x'}{y'}.$$

The equation  $yy' + xx' = c^2$

$$\text{gives } y = -\frac{x'}{y'}x + \frac{c^2}{y'}; \quad (1.)$$

and the equation  $x'^2 + y'^2 = c^2$

$$\text{gives } \frac{x'^2}{y'^2} + 1 = \frac{c^2}{y'^2}, \text{ or, } \frac{c^2}{y'} = c\sqrt{\frac{x'^2}{y'^2} + 1};$$

whence, by substitution in (1), we have

$$y = -\frac{x'}{y'}x + c\sqrt{\frac{x'^2}{y'^2} + 1};$$

or, writing  $m$  for  $-\frac{x'}{y'}$ ,

$$y = mx + c\sqrt{1 + m^2}; \quad (2.)$$

which is the equation to the tangent in terms of the tangent of the angle which the line makes with the axis of  $x$ .

Hence, if the general equation to a straight line be

$$y = mx + n,$$

then, to determine the condition that any straight line should touch the circle, we have

$$mx + n = mx + c\sqrt{1 + m^2},$$

$$\text{or, } n^2 = c^2(1 + m^2).$$

**30.** To find the equation to the normal at any point on the circumference of a circle.

The normal to a curve is the line intersecting the tangent at right angles at the point of contact.

Let  $(x', y')$  be the point of contact of a tangent to a circle. The equation to the tangent is

$$y = -\frac{x'}{y'}x + \frac{c^2}{y'}.$$

The equation to the normal as a straight line passing through  $(x', y')$  is of the form.

$$y - y' = m'(x - x');$$

and the condition of its being perpendicular to the tangent

requires that  $m'$  shall denote the reciprocal, with contrary sign, of the  $m$  found in the equation to the tangent; or requires that  $m' = -\frac{1}{m} = \frac{y'}{x'}$ ; thus the equation to the normal is

$$y - y' = \frac{y'}{x'}(x - x'),$$

or by reduction,  $x'y = xy'$ , or  $y = \frac{y'}{x'}x$ ;

which is the equation to a straight line passing through the origin, that is, through the centre.

The equation to the normal is anticipated in art. 28; where it is shewn that a tangent to a circle is always perpendicular to the radius passing through the point of contact.

**31.** *To find the locus of the middle points of a system of parallel chords.*

Let  $PP'$  be one of the chords;  $x', y'$  the coordinates of  $P$ ; and  $x'', y''$  those of  $P'$ .

The equation to  $PP'$  is

$$y - y' = \frac{y'' - y'}{x'' - x'}(x - x').$$

Now, because  $P$  and  $P'$  are points on the circumference,

$$c^2 = x'^2 + y'^2 = x''^2 + y''^2;$$


$$\therefore x''^2 - x'^2 = y'^2 - y''^2,$$

$$\text{or, } (x'' + x')(x'' - x') = (y' + y'')(y' - y'');$$

$$\therefore \frac{y'' + y'}{x'' + x'} = -\frac{x'' - x'}{y'' - y'}.$$

But the coordinates of  $V$ , the middle point of  $PP'$ , are  $\frac{1}{2}(x'' + x')$  and  $\frac{1}{2}(y'' + y')$ ; so that the tangent of the angle which a line through that point and the origin makes with the axis of  $x$  is

$$\frac{y'' + y'}{x'' + x'} = -\frac{x'' - x'}{y'' - y'}$$



=the reciprocal, with contrary sign, of the tangent number in the equation to  $PP'$ ; hence, the locus of  $V$  is a line through the centre perpendicular to the chords (*See Euclid, III. 3*).

**32.** *To find the equation to a straight line that shall touch a given circle, and pass through a given point external to the circle.*

Let  $h, k$  be the coordinates of the given point; and let  $x', y'$  (which will presently become known) be the coordinates of the point of contact.

The equation to the tangent at  $(x', y')$  is

$$xx' + yy' = c^2;$$

and, as the tangent passes through the given point  $(h, k)$ , the equation to the tangent must be satisfied by the coordinates of that point; and, therefore,

$$hx' + ky' = c^2; \text{ which is the required equation;}$$

but the point of contact being on the circumference, we have

$$x'^2 + y'^2 = c^2;$$

so that  $x'$  and  $y'$  are now determined, and will each have two values; therefore *two* tangents can be drawn to the circle from the given point; and

$$hx + ky = c^2$$

will be the general equation to the line joining the two points of contact, that is, to the *chord of contact*, as that line is called.

**33.** We shall now subjoin a few **Examples and Exercises** on articles **23–32**.

#### EXAMPLES.

*Ex. (1).* Find the points of intersection of the circle  $x^2 + y^2 = 4$  with the straight line  $3x + 5y = 6$ .

The 2nd equation gives  $9x^2 = 36 - 60y + 25y^2$ .

The 1st equation gives  $9x^2 = 36 - 9y^2$ .

By subtraction,  $34y^2 - 60y = 0$ ;

hence,  $y = 0$ , or  $\frac{3}{2}$ ;

$x = 2 - \frac{5}{2}y = 2$ , or  $-\frac{1}{2}$ .

Therefore the straight line meets the circle at the points  $(2, 0)$  and  $(-\frac{1}{2}, \frac{3}{2})$ ; the former of these points being an extremity of the diameter in the axis of  $x$ .

*Ex. (2).* Find the points of intersection of the circle  $x^2 + y^2 - 26x - 4y + 73 = 0$  with the line  $12x + 5y = 216$ .

The 2nd eqn. gives  $25y^2 = 144x^2 - 5184x + 46656$ .

„ 1st „  $25y^2 = -25x^2 + 650x - 1825 + 100y$ .

By subtraction,  $169x^2 - 5834x + 48481 = 100y$ ,  
that is, from 2nd eqn.  $= 4320 - 240x$ ;

hence,  $169x^2 - 5594x = -44161$ ;

which gives  $13x - 21\frac{2}{3} = \pm 6\frac{2}{3}$ .

Accordingly, we have  $x = 13$ , or  $20\frac{1}{3}$ ;

$y = 12$ , or  $-5\frac{1}{3}$ ;

and, therefore, the straight line meets the circle at the points  $(13, 12)$  and  $(20\frac{1}{3}, -5\frac{1}{3})$ ; the former of these being an extremity of the diameter parallel to the axis of  $y$ , since the abscissa, 13, is that of the centre, the equation to the circle being

$$(x-13)^2 + (y-2)^2 = 10^2.$$

*Ex. (3).* Find the length of the chord made by the circle  $x^2 + y^2 = c^2$  with the line  $\frac{x}{a} + \frac{y}{b} = 1$ .


Let the intercepts on the axes be  $OA = a$ ,  $OB = b$ ,  $\therefore AB^2 = a^2 + b^2$ ; and  $OR$ , the perpendicular on  $AB$ , is determined, by similar triangles, thus:

$$AB^2 : OB^2 :: OA^2 : OR^2 = \frac{a^2 b^2}{a^2 + b^2};$$

Now,  $OR$  bisects the chord, and

$$\text{therefore half the chord} = \sqrt{c^2 - \frac{a^2 b^2}{a^2 + b^2}};$$

$$\therefore \text{the length of the chord is} = 2\sqrt{c^2 - \frac{a^2 b^2}{a^2 + b^2}}.$$



*Ex. (4).* Find the equation to the circle described on a diameter of which the extremities are  $(x', y')$  and  $(x'', y'')$ .

Here the coordinates of the centre are  $\frac{1}{2}(x' + x'')$  and  $\frac{1}{2}(y' + y'')$ , therefore,

$$\left\{ x - \frac{1}{2}(x' + x'') \right\}^2 + \left\{ y - \frac{1}{2}(y' + y'') \right\}^2 = \text{rad.}^2$$


$$= \frac{1}{4}(x'' - x')^2 + \frac{1}{4}(y'' - y')^2;$$

or  $x^2 - (x' + x'')x + \frac{1}{4}(x'' + x')^2 + y^2 - (y' + y'')y + \frac{1}{4}(y'' + y')^2$   
 $- \frac{1}{4}(x'' - x')^2 - \frac{1}{4}(y'' - y')^2 = 0$

or,  $x^2 + y^2 - (x' + x'')x - (y' + y'')y + x'x'' + y'y'' = 0$ ; the equation required.

*Ex. (5).* Given AB the base of a triangle, and the ratio of its sides AP, BP =  $m$ : shew that the locus of the vertex P is a circle, except when  $m=1$ .

Take the base AB as the axis of  $x$ , and O the middle point of AB as origin. Let  $x, y$  be the coordinates of the vertex P; and put  $AB=2a$ .



If  $AP=m$  BP, then  $AP^2=m^2BP^2$ ;

$$\text{or, } AM^2 + PM^2 = m^2(BM^2 + PM^2);$$

$$\text{that is, } (a+x)^2 + y^2 = m^2 \left\{ (a-x)^2 + y^2 \right\};$$

$$\text{or } m^2(x^2 + y^2 - 2ax + a^2) = x^2 + y^2 + 2ax + a^2;$$

that is,


$$(m^2-1)x^2 + (m^2-1)y^2 - 2(m^2+1)ax + (m^2-1)a^2 = 0;$$

$$\text{hence, } x^2 + y^2 - 2 \frac{m^2+1}{m^2-1} ax + a^2 = 0;$$

which is the equation to a circle, unless  $m=1$ ; for when  $m=1$  the equation becomes  $x^2 + y^2 - \infty + a^2 = 0$ ; then we should have  $x=0$ , and  $y$  indefinite or arbitrary, and  $BP^2 = y^2 + a^2 = AP^2$ ; in which case the locus would be the axis of  $y$ .

*Ex. (6).* In one plane, the locus of the point from which two given unequal circles would appear equally large, that is, subtend equal angles, is a circle.

Let  $AC$  the radius of the smaller circle  $=r$ ,  $BD$  that of the larger  $=r'$ , and  $AB$  the distance of the centres  $=a$ . Let  $P$  be a point from which the circles subtend equal angles, viz. doubles of  $APC$ ,  $BPD$ ;  $PC$  and  $PD$  being tangents.



Draw  $EF$  a common tangent to the circles, and let it meet  $AB$  at  $O$ . Take  $O$  as origin,  $x, y$  as the coordinates of  $P$ .

By similar triangles,  $\frac{OA}{OB-OA} = \frac{AE}{BF-AE}$ ; hence, if we

put  $d$  for  $OA$ , we have  $\frac{d}{a} = \frac{r}{r'-r}$ , or  $d = \frac{ar}{r'-r}$ .

$$AM = x - d, \quad BM = a + d - x, \quad PM = y.$$

By similar triangles,  $\frac{PA^2}{PB^2} = \frac{AC^2}{BD^2} = \frac{r^2}{r'^2}$ ;

$$\text{that is, } \frac{(x-d)^2 + y^2}{(d+a-x)^2 + y^2} = \frac{r^2}{r'^2};$$

$$\text{whence } (x^2 + y^2)(r'^2 - r^2) - 2x\{r'^2d - r^2(d+a)\} \\ = r^2(d+a)^2 - r'^2d^2, = 0, \text{ since } d+a = \frac{ar'}{r'-r};$$

$$\text{moreover, } r'^2d - r^2(d+a) = \frac{arr'^2 - ar'^2r}{r'-r} = arr';$$

$$\therefore x^2 + y^2 - 2x \cdot \frac{arr'}{r'^2 - r^2} = 0,$$

which is the equation to a circle, whose centre is on the axis of  $x$  at a point  $Q$  equidistant from the points  $O$  and  $P$ ;

the radius  $OQ = PQ = \frac{arr'}{r'^2 - r^2}$ .

## EXERCISES [C].

1. Determine the radius, and the coordinates of the centre, for each of the following circles :

(i.)  $x^2 + y^2 - 6x - 8y - 24 = 0$ .

(ii.)  $x^2 + y^2 + 2x + 4y - 4 = 0$ .

(iii.)  $x^2 + y^2 - 6x - 18y + 10 = 0$ .

(iv.)  $x^2 + y^2 + x - 2y - 3\frac{3}{4} = 0$ .

2. Find the points of intersection of the circle  $x^2 + y^2 = 34$  :

(i.) with the line  $x - y = 2$ .

(ii.) with the line  $2x + y = -1$ .

3. Find the points of intersection of the circle  $x^2 + y^2 = 1$  :

(i.) with the line  $3x - 4y = 5$ .

(ii.) with the line  $5x + 4y = -5$ .

4. Find the points of intersection of the line

$$4y - 3x + 32 = 0 :$$

(i.) with the circle  $x^2 + y^2 - 24x - 10y = 0$ .

(ii.) with the circle  $2x^2 + 2y^2 - 8x + y - 45 = 0$ .

5. A straight line parallel to the line  $y = mx + n$  touches the circle  $(x-a)^2 + (y-b)^2 = c^2$ . Find the equation to the tangent.

6. Find the length of the chord which the line

$$12y - 5x = 56$$

makes with the circle  $x^2 + y^2 - 40x = 0$ .

7. What must be the relation between the quantities

$a$ ,  $b$ ,  $c$ , in order that the line  $\frac{x}{a} + \frac{y}{b} = 1$  may touch the circle  $x^2 + y^2 = c^2$ ?

8. Two circles, whose radii are  $c$ ,  $c'$ , cut one another; the distance between their centres being  $a$ . Find the coordinates of the points of intersection.

9. From the equation to the circle, which is the locus of P in the 5th of the worked Examples, viz.,

$$x^2 + y^2 - 2 \cdot \frac{m^2 + 1}{m^2 - 1} ax + a^2,$$

determine the position of the centre, the length of the radius, and the points at which the circle cuts the axis of  $x$ .

10. Shew that the square of half the base of a triangle and the square of the line joining the vertex and the middle of the base are together equal to half the sum of the squares of the two sides of the triangle. Shew also that, when the base and the sum of the squares of the sides are constant, the locus of the vertex is a circle.

11. Find the equation to a circle whose centre is at the origin of coordinates, and which touches the line  $y=3x+5$ .

12. Find the equation to the circle whose diameter is the common chord of the circles  $x^2+y^2=c^2$  and  $(x-a)^2+y^2=c^2$ , where  $a$  denotes the distance between the two centres on the axis of  $x$ .

13. What must be value of  $c$ , in order that the circles  $(x-a)^2+(y-b)^2=c^2$  and  $(x-b)^2+(y-a)^2=c^2$  may touch each other?


14. Suppose the equation to a circle whose radius is  $c$  to be  $x^2+y^2-2cx=0$ , and the straight line joining its points of intersection with the line  $y=mx$  to have a circle described on it as diameter. Required the equation to the latter circle.



## THE ELLIPSE.

**34.** The Ellipse is the locus of a point whose distance from a given fixed point is always less, in a constant ratio, than its distance from a given fixed straight line.

Let  $S$  be the given point, and  $KK'$  the given line. Draw



$SD$  perpendicular to  $KK'$ . Let  $P$  be a point on the locus; join  $SP$ , and draw  $PM$  parallel to  $KK'$  and  $PN$  parallel to  $DS$ .

Take  $D$  as the origin,  $DS$  as the axis of  $x$ ,  $KK'$  as the axis of  $y$ .

Divide  $DS$  at  $A'$  so that  $\frac{A'S}{A'D} = \frac{SP}{PN}$  = the constant ratio, a value less than unity, which we may call  $e$ . Let  $DS$  be called  $p$ ; and let  $x, y$  be the coordinates of  $P$ .

We have  $SP = e \cdot PN$ ;  $\therefore SP^2 = e^2 PN^2$ ; or  $PM^2 + SM^2 = e^2 DM^2$ ; that is,  $y^2 + (x-p)^2 = e^2 x^2$ , which is the equation to the ellipse with the assumed origin and axes.

**35.** For the point where the ellipse meets the axis of  $x$ , we have  $y=0$ , and the equation becomes

$$(x-p)^2 = e^2 x^2;$$

but this gives  $x-p = \pm ex$ , or  $(1 \mp e)x = p$ , or  $x = \frac{p}{1 \mp e}$ ;

signifying that the ellipse meets the axis at two points,  $A'$  and  $A$ ; so that  $DA' = \frac{p}{1+e}$ , and  $DA = \frac{p}{1-e}$ .

$C$  the middle point of  $A'A$  is called the centre, and the extremities  $A'$ ,  $A$  are called the vertices of the ellipse. Also, the fixed point  $S$  is called the focus, and the fixed line  $KK'$  the directrix of the ellipse.

**36.** Let the line  $A'A = 2a$ ; then  $A'C = a$ .

Since  $A$  is a point on the locus,

$$\text{we have } e = \frac{SA}{DA} = \frac{2a - A'S}{2a + DA'} = \frac{2a - e \cdot DA'}{2a + DA'};$$

$$\therefore a = ae + e \cdot DA'.$$

Whence are derived the following values :

$$(i.) DA' = \frac{a}{e}(1-e);$$

$$(ii.) A'S = e \cdot DA' = a(1-e);$$

$$(iii.) DC = a + DA' = \frac{a}{e};$$

$$(iv.) SC = a - A'S = ae;$$

$$(v.) DS = p = DC - SC = \frac{a}{e} - ae = \frac{a}{e}(1-e^2).$$

**37.** Suppose the origin to be at  $A'$ .

In this case  $A'M$  takes the place of  $DM$ ; and if  $A'M$  be called  $x'$ , then  $DM = DA' + x'$ ; that is,  $x = x' + \frac{a}{e} - a$ ; whence, by substitution in the equation  $y^2 + (x-p)^2 = e^2x^2$ , we have

$$y^2 + \left(x' + \frac{a}{e} - a - \frac{a}{e} + ae\right)^2 = \{ex' + a(1-e)\}^2;$$

$$\text{or } y^2 + \{x' - a(1-e)\}^2 = \{ex' + a(1-e)\}^2;$$

$$\text{or } y^2 + x'^2 - 2a(1-e)x' = e^2x'^2 + 2ae(1-e)x';$$

$$\text{or } y^2 + x'^2 - 2a(1+e)(1-e)x' = e^2x'^2;$$

$$\text{or } y^2 + x'^2(1-e^2) = 2ax'(1-e^2);$$

$$\text{or } y^2 = (1-e^2)(2ax' - x'^2).$$

Now, as A is to be accounted the origin, we may remove the accent, and the equation will be

$$y^2 = (1 - e^2)(2ax - x^2).$$

**38.** Suppose the origin to be at C.

Here again we might put  $x'$  for CM, and substitute  $x' + \frac{a}{e}$  for  $x$  in the original equation.

Or we may at once let  $x$  and  $y$  be the coordinates; then we have

$$SM = CM + SC = x + ae;$$

$$DM = CM + DC = x + \frac{a}{e},$$

$$\text{or } e \cdot DM = ex + a;$$

$\therefore$ , equivalent to  $PM^2 + SM^2 = e^2 DM^2$ , we have

$$y^2 + (x + ae)^2 = (ex + a)^2;$$

$$\text{or } y^2 = (1 - e^2)(a^2 - x^2).$$

Now, let  $b$  denote the ordinate BC; then for the coordinates of the point B we have  $x=0$ ,  $y=b$ , and the above equation gives us  $b^2 = a^2(1 - e^2)$ , or  $1 - e^2 = \frac{b^2}{a^2}$ . The equation, therefore, may be written

$$y^2 = \frac{b^2}{a^2}(a^2 - x^2),$$

which is reducible to the more symmetrical form

$$a^2 y^2 + b^2 x^2 = a^2 b^2, \text{ or } \frac{x^2}{a^2} + \frac{y^2}{b^2} = 1;$$

which is the equation to the ellipse referred to the centre as origin, and is the one most frequently employed.

**39.** The above equation is convertible into the forms

$$y = \pm \frac{b}{a} \sqrt{a^2 - x^2}, \quad (1)$$

$$\text{and } x = \pm \frac{a}{b} \sqrt{b^2 - y^2}, \quad (2)$$

which enable us readily to determine the form of the ellipse.

If  $y=0$ , then (2) gives  $x=\pm a$ ; therefore the ellipse cuts the axis of  $x$  at the points  $A, A'$ .

If  $x$  be supposed a greater magnitude than  $+a$  or  $-a$ , then in (1) we shall have  $a^2-x^2$  negative, and the values of  $y$  are imaginary. Therefore no point on the ellipse can have a greater abscissa than  $a$  or  $-a$ .

If  $x$  be supposed a less magnitude than  $+a$  or  $-a$ , then (1) shews that for every value of  $x$  there are two values of  $y$ , numerically equal, but contrary in sign. This implies that for every point above the axis of  $x$  there is a corresponding point at the same distance below the axis of  $x$ ; so that the axis of  $x$  divides the curve symmetrically.

Also from (2) it is evident that the curve is symmetrical with respect to the axis of  $y$ , and that no point on the ellipse has an ordinate exceeding  $b$ .

Accordingly, the ellipse is a continuous curve returning into itself, and divided by the axis of  $x$  into two equal parts; and it is easily shewn that *the centre of an ellipse is a point that bisects every chord of the ellipse drawn through that point.*

It may be here stated that the term *axes*, in reference to the ellipse, is generally used to denote the lines  $A'A$  and  $BB'$  intercepted by the curve; and that the former is called the major axis or transverse axis, the latter the minor or conjugate axis.

It may be also mentioned that the ratio denoted by the symbol  $e$  is called the *eccentricity* of the ellipse.

**40.** The ellipse being symmetrically situated with respect to the axes,  $A'A, BB'$ , there must be two foci, each having its corresponding directrix; and if  $AH=A'S$ , and  $AE=DA'$ , the curve can be described by means of  $H$  and  $LL'$ , as focus and directrix.

**41.** The readiest mechanical method of describing an ellipse is furnished by the following property:

The sum of the focal distances of any point on the ellipse is equal to the axis major.

$$\text{For } SP = e \cdot PN = e(DC + CM) = e\left(\frac{a}{e} + x\right) = a + ex;$$

$$\text{and } HP = e \cdot PR = e(CE - CM) = e\left(\frac{a}{e} - x\right) = a - ex;$$

$$\therefore SP + PH = 2a = A'A.$$

Hence, for the point B we have  $SB = A'C$ .

Accordingly, an ellipse may be constructed as follows:

Let a string SPH be fastened to two points S and H, then, if a pencil P be moved so as to keep the string always stretched, it will trace an ellipse of which S and H are the foci, since  $SP + PH$  will be constantly  $= A'A$ .

This property of the constant sum of the focal distances is sometimes stated as a definition of the ellipse (*See Ex. 4, p. 54*).

**42.** Either focus divides the axis major into segments, the rectangle of which is equal to the square of the semi-axis minor, so that  $A'S \cdot SA = BC^2$ .

For  $A'C^2 = SB^2 = SC^2 + BC^2$ ; therefore  $BC^2 = A'C^2 - SC^2 = (A'C - SC)(A'C + SC) = A'S \cdot SA$ .

Or thus:  $A'S = a(1 - e)$ , and  $SA = a(1 + e)$ ; and we have already found (38) the product of these, viz.  $a^2(1 - e^2) = b^2$ .

**43.** The equation  $y^2 = \frac{b^2}{a^2}(a^2 - x^2)$  may be put in the form

$$y^2 = \frac{b^2}{a^2}(a + x)(a - x);$$

hence, if PM be any ordinate, and C the origin,

$$PM^2 = \frac{BC^2}{A'C^2} A'M \cdot MA,$$

$$\text{or } PM^2 : A'M \cdot MA :: BC^2 : A'C^2.$$

**44.** The double ordinate FHF' passing through the focus is called the *latus rectum*. To find its length, we have  $x = CH = SC = ae$ , which substituted for  $x$  in the equation

$y^2 = \frac{b^2}{a^2}(a^2 - x^2)$  gives

$$y^2 = \frac{b^2}{a^2}(a^2 - a^2e^2) = b^2(1 - e^2) = b^2 \cdot \frac{b^2}{a^2} = \frac{b^4}{a^2};$$

$$\text{hence } y = FH = \frac{b^2}{a}; \text{ and } FHF' = \frac{2b^2}{a}.$$

#### TANGENT AND NORMAL TO AN ELLIPSE.

**45.** The tangent to an ellipse may be regarded as a secant passing through two coincident points on the curve. (See art. 27.)

*To find the equation to the tangent at a given point on an ellipse.*

Let  $x, y$  be the coordinates of P the given point,  $x', y'$  those of another point P' on the curve, near the given point.

The equation to the straight line PP' is

$$y - y' = \frac{y'' - y'}{x'' - x'}(x - x'). \quad (1)$$

The equation to the ellipse, referred to the centre as origin, is

$$a^2y^2 + b^2x^2 = a^2b^2.$$

Accordingly, since  $(x', y')$  and  $(x'', y'')$  are points on the curve

$$\begin{aligned} a^2y'^2 + b^2x'^2 &= a^2b^2, \\ a^2y''^2 + b^2x''^2 &= a^2b^2; \\ \therefore a^2(y''^2 - y'^2) &= -b^2(x''^2 - x'^2), \\ \text{or, } a^2(y'' + y')(y'' - y') &= -b^2(x'' + x')(x'' - x'); \\ \therefore \frac{y'' - y'}{x'' - x'} &= -\frac{b^2}{a^2} \cdot \frac{x'' + x'}{y'' + y'}; \end{aligned}$$

hence, the 1st equation, by substitution, will become

$$y - y' = -\frac{b^2}{a^2} \cdot \frac{x'' + x'}{y'' + y'}(x - x'). \quad (2)$$

Now, when the point  $(x'', y'')$  coincides with the point  $(x', y')$ , we have  $x'' = x'$  and  $y'' = y'$ ; in which case (2)

$$\text{becomes} \quad y - y' = -\frac{b^2}{a^2} \cdot \frac{2x'}{2y'}(x - x'),$$

that is, 
$$y - y' = -\frac{b^2 x'}{a^2 y'} (x - x'),$$
or  $a^2 y y' + b^2 x x' = a^2 y'^2 + b^2 x'^2,$ 
that is,  $a^2 y y' + b^2 x x' = a^2 b^2.$  (3)

**46.** To express the tangent to an ellipse in terms of the tangent of the angle which the line makes with the axis of  $x$ .

We have seen that the tangent of the angle which the line makes with the major axis is  $-\frac{b^2 x'}{a^2 y'}.$

From the equation  $a^2 y y' + b^2 x x' = a^2 b^2$   
we have 
$$y = -\frac{b^2 x'}{a^2 y'} x + \frac{b^2}{y'}; \quad (1)$$

and from the equation  $a^2 y'^2 + b^2 x'^2 = a^2 b^2$   
we have 
$$1 + \frac{b^2 x'^2}{a^2 y'^2} = \frac{b^2}{y'^2};$$

$$\therefore \frac{b^2}{y'} = \pm b \sqrt{\frac{b^2 x'^2}{a^2 y'^2} + 1}. \quad (2)$$

Let  $-\frac{b^2 x'}{a^2 y'} = m$ ; whence  $\frac{x'}{y'} = -\frac{a^2 m}{b^2};$

$$\therefore \frac{x'^2}{y'^2} = \frac{a^4 m^2}{b^4}; \text{ and } \frac{b^2 x'^2}{a^2 y'^2} = \frac{a^2 m^2}{b^2};$$

hence, by substitution in (2),

$$\frac{b^2}{y'} = \pm b \sqrt{\frac{a^2 m^2}{b^2} + 1}; \text{ or, } \frac{b^2}{y'} = \pm \sqrt{a^2 m^2 + b^2}.$$

Whence, by substitution in (1),

$$y = mx \pm \sqrt{a^2 m^2 + b^2}; \quad (3)$$

which is the equation to the tangent in terms of the tangent of the angle which the line makes with the major axis of the ellipse; the double sign implying that *two* tangents may be drawn making angles of equal magnitude with the axis of  $x$ . Hence, if the general equation to a straight line be

$$y = mx + n,$$

then, to determine the condition that any straight line should touch the ellipse, we have

$$mx + n = mx + \sqrt{a^2m^2 + b^2},$$

$$\text{or } n^2 = a^2m^2 + b^2.$$

**47.** To find the equation to the normal at any point of an ellipse, and its intercepts on the axes.


The *normal*, as formerly defined, is the line intersecting the tangent at right angles at the point of contact.

Let  $(x', y')$  be the point of contact of a tangent to the ellipse. The equation to the tangent is

$$y = -\frac{b^2x'}{a^2y'}x + \frac{b^2}{y'}.$$

The equation to the normal as a straight line through  $(x', y')$  is of the form  $y - y' = m'(x - x')$ ; and the condition of its being perpendicular to the tangent requires that  $m'$  shall denote the reciprocal, with contrary sign, of the  $m$  found in the equation to the tangent, or requires that  $m' = -\frac{1}{m} = \frac{a^2y'}{b^2x'}$ ; hence the equation to the normal at  $(x', y')$  is

$$y - y' = \frac{a^2y'}{b^2x'}(x - x').$$



Now, at G, where the normal PG cuts the axis major,  $y=0$ , and the above equation gives

$$-y' = \frac{a^2y'}{b^2x'}(x - x'); \text{ or, } -1 = \frac{a^2}{b^2x'}(x - x');$$



$$\text{or, } x = x' \left( 1 - \frac{b^2}{a^2} \right) = e^2 x';$$

that is,  $CG = e^2 CM$ .

Again, at  $G'$ , where the normal cuts the axis minor,  $x=0$ , and the equation to the normal gives

$$y - y' = \frac{a^2 y'}{b^2 x'} (0 - x'); \text{ or, } y = y' - \frac{a^2 y'}{b^2};$$

$$\text{or, } y = y' \left( -1 \frac{a^2}{b^2} \right) = \frac{a^2 e^2}{b^2} y';$$

that is,  $CG' = \frac{a^2 e^2}{b^2} PM$ .

*Note.* The length of  $PG$  in terms of the focal distances of  $P$  and the eccentricity may be determined as follows:—

$$SP = a + ex'; \quad HP = a - ex'; \quad SM = ae + x'.$$

$$\therefore PM^2 = SP^2 - SM^2 = (a^2 - x'^2)(1 - e^2);$$

$$GM = CM - CG = x'(1 - e^2);$$

$$GM^2 = x'^2(1 - e^2)^2.$$

$$\text{Hence } PG^2 = PM^2 + GM^2 = (1 - e^2) \{ a^2 - [1 - (1 - e^2)] x'^2 \}$$

$$= (1 - e^2) (a^2 - e^2 x'^2) = (1 - e^2) (a + ex')(a - ex');$$

$$\text{or } PG^2 = (1 - e^2) SP \cdot HP.$$

**48.** *The focal distances of any point on the ellipse make equal angles with the tangent at that point.*

$C$  being the origin,  $SP$  and  $HP$  the focal distances of  $P$ , and  $SC = CH = ae$ ; the coordinates of  $S$  are  $-ae, 0$ , and those of  $H$  are  $ae, 0$ : let those of  $P$  be  $x', y'$ .

The equation to  $SP$  is

$$y - 0 = \frac{y' - 0}{x' + ae} (x + ae),$$

showing the value of  $\tan \angle PSM$  to be  $\frac{y'}{x' + ae}$ .

The equation to  $HP$  is

$$y - 0 = \frac{y' - 0}{x' - ae} (x - ae),$$

showing the value of  $\tan \text{PHT}$  to be  $\frac{y'}{x' - ae}$  ;

$\therefore$  that of  $\tan \text{PHM}$  is  $-\frac{y'}{x' - ae}$ .

The equation to the tangent at P is

$$y - y' = -\frac{b^2 x'}{a^2 y'} (x - x'),$$

showing the value of  $\tan \text{PTM}$  to be  $\frac{b^2 x'}{a^2 y'}$ .

$\tan \text{SPT}' = \tan(\text{PSM} + \text{PTM})$

$$\begin{aligned} &= \frac{\frac{y'}{x' + ae} + \frac{b^2 x'}{a^2 y'}}{1 - \frac{b^2 x'}{a^2 (x' + ae)}} = \frac{a^2 y'^2 + b^2 x'^2 + ab^2 ex'}{(a^2 - b^2) x' y' + a^3 ey'} \\ &= \frac{a^2 b^2 + ab^2 ex'}{a^2 e^2 x' y' + a^3 ey'} = \frac{b^2 (a + ex')}{aey' (a + ex')} = \frac{b^2}{aey'}. \end{aligned}$$

$\tan \text{HPT} = \tan (\text{PHM} - \text{PTM})$

$$\begin{aligned} &= \frac{-\frac{y'}{x' - ae} - \frac{b^2 x'}{a^2 y'}}{1 - \frac{b^2 x'}{a^2 (x' - ae)}} = \frac{a^2 y'^2 + b^2 x'^2 - ab^2 ex'}{(b^2 - a^2) x' y' + a^3 ey'} \\ &= \frac{a^2 b^2 - ab^2 ex'}{-a^2 e^2 x' y' + a^3 ey'} = \frac{b^2 (a - ex')}{aey' (a - ex')} = \frac{b^2}{aey'}. \end{aligned}$$

Therefore the angle  $\text{SPT}' = \text{the angle HPT}$ . And because PG is perpendicular to TTV we have also the complements SPG, HPG equal to each other ; so that the angle between the focal distances of any point on the ellipse is bisected by the normal at that point.

The above is a rigorous analytical proof of the proposed theorem. The following is another demonstration, assuming the 2nd part of Euclid VI. 3.

In (47) we found  $\text{CG} = e^2 x'$  ;

$\therefore \text{SG} = \text{SC} + \text{CG} = ae + e^2 x' = e(a + ex') = e \cdot \text{SP}$  (art. 41)

$$HG = CH - CG = ae - e^2x' = e(a - ex') = e \cdot HP \quad (\text{art. 41})$$

$$\therefore \frac{SG}{HG} = \frac{SP}{HP};$$


hence (Eucl. VI. 3)  $SPG = HPG$ ; and  $\therefore SPT' = HPT$ .

**49.** If a circle be described on the axis major as diameter, and any ordinate,  $MP$ , of the ellipse be produced to meet the circle at  $P'$ , then

$$\frac{PM}{P'M} = \frac{b}{a}.$$

For  $CM^2 + P'M^2 = CA^2 = a^2$ , or,  $a^2 - CM^2 = P'M^2$ ; and the equation to the ellipse gives  $\frac{CM^2}{a^2} + \frac{PM^2}{b^2} = 1$ , or,  $a^2 - CM^2 = \frac{a^2}{b^2} PM^2$ . Therefore  $\frac{a^2}{b^2} PM^2 = P'M^2$ ,

$$\text{or } \frac{PM}{P'M} = \frac{b}{a}.$$



The angle  $P'CM$  is called the *eccentric angle* of the ellipse.

**50.** To find the locus of the intersection of a tangent to the ellipse with the perpendicular drawn to it from either focus.

The equation to  $RR'$  (see figure in 49) is

$$y - mx = \sqrt{a^2m^2 + b^2}. \quad (1)$$

The equation to  $SR'$  drawn from the focus whose coordinates are  $-ae, 0$ , is of the form

$$y = m'(x + ae);$$

and that this line may be perpendicular to the former we put  $m' = -\frac{1}{m}$ , or

$$y = -\frac{1}{m}(x + ae);$$

$$\text{or, } my + x = -ae. \quad (2)$$

Similarly, the equation to  $HR$  drawn perpendicular from the focus whose coordinates are  $ae, 0$ , will be

$$y = -\frac{1}{m}(x - ae),$$

$$\text{or } my + x = ae. \quad (3)$$

Squaring (1) and (2), or (1) and (3):

$$y^2 - 2myx + m^2x^2 = a^2m^2 + b^2;$$

$$m^2y^2 + 2myx + x^2 = a^2e^2.$$

By addition

$$y^2(1 + m^2) + x^2(1 + m^2) = a^2m^2 + b^2 + a^2e^2$$

$$= a^2m^2 + a^2 = a^2(1 + m^2);$$


$$\therefore y^2 + x^2 = a^2,$$

the equation to a circle whose radius is  $a$ . Therefore the required locus is a circle of which the major axis of the ellipse is a diameter.

*Note.* The preceding result suggests the following geometrical method of drawing a tangent to an ellipse from any external point  $Q$ .

Let a circle be described on the major axis as diameter, and another on the line  $SQ$  as diameter; then straight lines, drawn from

$Q$  through  $R, R'$ , the points of intersection of the two circles, will be the two tangents to the ellipse.



For the angles QRS, QR'S in a semicircle are right angles, or R, R' are points on the locus of the intersection of a perpendicular from the focus with the tangent.

**51.** To determine the perpendicular ( $p$ ) from the centre of an ellipse as origin, to the tangent at any point on the curve, in terms of the angle  $\theta$  which  $p$  makes with the axis of  $x$ .

(See the fig. in art. 49.)  $CV=p$ ;  $VCT=\theta$ .

The equation to the straight line TV is

$$\frac{x}{a} + \frac{y}{b} = 1;$$

multiplying by  $p$ , we have

$$\frac{p}{a}x + \frac{p}{b}y = p; \text{ that is, } \frac{CV}{CT}x + \frac{CV}{CI}y = p;$$

but  $\frac{CV}{CT} = \cos \theta$ , and  $\frac{CV}{CI} = \sin \theta$ ;

$$\therefore x \cos \theta + y \sin \theta = p;$$

which, it should be remembered, is a convenient expression for a straight line in terms of the perpendicular on it from the origin and the inclination of this perpendicular to the axis of  $x$ . It gives

$$y = -\frac{\cos \theta}{\sin \theta}x + \frac{p}{\sin \theta};$$

and the condition that this line should touch the ellipse is

$$\frac{p^2}{\sin^2 \theta} = a^2 \frac{\cos^2 \theta}{\sin^2 \theta} + b^2; \text{ or } p^2 = a^2 \cos^2 \theta + b^2 \sin^2 \theta;$$

$$\text{or } p^2 = a^2 \cos^2 \theta + a^2(1-e^2)\sin^2 \theta;$$

$$= a^2(\cos^2 \theta + \sin^2 \theta) - a^2 e^2 \sin^2 \theta;$$

$$\text{that is, } p^2 = a^2(1 - e^2 \sin^2 \theta);$$

which determines  $p$  as required.

**52.** In this article we shall give a few Examples and Exercises on the Ellipse.

EXAMPLES.

*Ex. (1).* Find the eccentricity of the ellipse

$$5x^2 + 8y^2 = 2n^2.$$

Here, in order to make the equation assume the form

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1,$$

we divide it by  $2n^2$ , which gives

$$\frac{x^2}{\frac{2}{5}n^2} + \frac{y^2}{\frac{1}{4}n^2} = 1;$$

$$\therefore a^2 = \frac{2}{5}n^2, b^2 = \frac{1}{4}n^2, \text{ and } \frac{b^2}{a^2} = \frac{5}{8}.$$

$$\text{Now } e^2 = 1 - \frac{b^2}{a^2} = 1 - \frac{5}{8} = \frac{3}{8} = \frac{1}{\frac{8}{3}} \text{ of } 6;$$

$$\therefore e = \frac{1}{2}\sqrt{6}.$$

*Ex. (2).* Find the eccentricity of the ellipse in which the normal at F, the extremity of the latus rectum, passes through B', the extremity of the axis minor.

$$\text{From art. 47 we have } CB' = \frac{a^2 e^2}{b^2} FH;$$

$$\text{that is, } b = \frac{a^2 e^2}{b^2} \cdot \frac{b^2}{a} = a e^2;$$

$$\therefore e^4 = \frac{b^2}{a^2} = 1 - e^2; \text{ or } e^4 + e^2 = 1;$$

$$\text{whence } e = \frac{1}{2}\sqrt{(2\sqrt{5}-2)}.$$

*Ex. (3).* The focal distances of a point P, on an ellipse, are SP=18, HP=8, and the semi-axis minor is 5. Find the coordinates of P referred to the centre as origin.

$$(\text{See the fig. in 34.}) \quad a = \frac{1}{2}(18+8) = 13; b = 5;$$

$$\text{hence } e = \sqrt{\left(1 - \frac{b^2}{a^2}\right)} = \frac{12}{13}; \text{ also } SC = ae = 12, SH = 24,$$

$$SM^2 - MH^2 = SP^2 - HP^2;$$

$$\text{that is, } (SM + MH)(SM - MH) = (SP + HP)(SP - HP),$$

$$\text{or } 24(\text{SM} - \text{MH}) = 26 \times 10; \therefore \text{SM} - \text{MH} = 10\frac{5}{6};$$

$$\text{hence SM} = 17\frac{5}{12};$$

$$\therefore \text{CM} = \text{SM} - \text{SC} = 5\frac{5}{12}.$$

$$\text{PM}^2 = (a^2 - x'^2)(1 - e^2) = (169 - \frac{4 \times 225}{144}) \times \frac{25}{169},$$

$$= 25 - \frac{25}{144} \text{ of } 25 = \frac{119}{144} \text{ of } 25 = \frac{25}{144} \text{ of } 119;$$

$$\therefore \text{PM} = \frac{5}{12} \sqrt{119};$$

therefore the required coordinates are  $x' = 5\frac{5}{12}$ ,  $y' = \frac{5}{12} \sqrt{119}$ .

*Ex. (4).* Investigate the equation to the curve that is the locus of a point, the sum of whose distances from two given points is constant.

Let S, H be the two fixed points, and take C the middle point of SH as origin. Let P be a point on the locus, and let

$$\text{SP} + \text{HP} = 2a, \text{ and } \text{SH} = 2c.$$

$$\text{SP}^2 = (c+x)^2 + y^2,$$

$$\text{HP}^2 = (c-x)^2 + y^2,$$

$$\therefore \text{SP}^2 - \text{HP}^2 = 4cx;$$

$$\frac{\text{SP}^2 - \text{HP}^2}{\text{SP} + \text{HP}} = \text{SP} - \text{HP} = \frac{4cx}{2a} = \frac{2cx}{a};$$

$$\text{hence, SP} = a + \frac{cx}{a}.$$


$$\therefore \left(a + \frac{cx}{a}\right)^2 = (c+x)^2 + y^2;$$

$$\text{which gives } a^2 + \frac{c^2 x^2}{a^2} = c^2 + x^2 + y^2;$$

$$\text{whence } y^2 = \frac{a^2 - c^2}{a^2} (a^2 - x^2);$$

which is the equation to an ellipse having  $a^2 - c^2 = b^2$ , that is, having S and H for the foci.

*Ex. (5).* Shew that the locus of one end of a given straight rod, whose other end and a given point in it move in straight lines at right angles to each other, is an ellipse.







but if  $m=t$  or  $t'$ , then  $m-t=0$ , and  $m-t'=0$ ,

$$\therefore (m-t)(m-t')=0,$$

$$\text{or } m^2-(t+t')m+tt'=0,$$

which compared with (1) shows  $tt'=\frac{b^2-k^2}{a^2-h^2}$ ;

$$\therefore \frac{b^2-k^2}{a^2-h^2}+1=0, \text{ or } b^2-k^2=h^2-a^2,$$

$$\text{or } h^2+k^2=a^2+b^2;$$

hence the locus of the point  $(h, k)$  is a circle.

*Ex. (7).* Find, in terms of the focal distances of a point P on the ellipse, the perpendiculars HR, SR' from the foci on the tangent at P; and shew that  $HR \cdot SR' = b^2$ .

(See the figures in arts. 47, 49.)

The equation to the tangent at P is

$$a^2yy'-b^2xx'=-a^2b^2.$$

For the point T, where it meets the axis of  $x$ , we have

$$\begin{aligned} y=0, \therefore x &= CT = \frac{a^2}{x'}; \text{ and } GT = CT - CG = \frac{a^2}{x'} - e^2x' \\ &= \frac{a^2 - e^2x'^2}{x'}; \text{ also } HT = GT - GH = \frac{a^2 - e^2x'^2}{x'} - e(a - ex) \\ &= \frac{a(a - ex')}{x'}. \end{aligned}$$

Now, by similar triangles,  $\frac{HR^2}{PG^2} = \frac{HT^2}{GT^2}$ ,

$$\text{or, } \frac{HR^2}{(1-e^2)(a^2-e^2x'^2)} = \frac{a^2}{(a+ex')^2}; \text{ or, } \frac{HR^2}{\frac{b^2}{a^2}(a-ex')} = \frac{a^2}{a+ex'}$$

$$\text{that is, } \frac{a^2}{b^2} \cdot \frac{HR^2}{HP} = \frac{a^2}{SP};$$

$$\text{hence, } HR^2 = \frac{HP}{SP} \cdot b^2.$$

Similarly, it may be shewn that the other perpendicular is determined by  $SR'^2 = \frac{SP}{HP} \cdot b^2$ .

$$\therefore HR \cdot SR' = b^2.$$


## EXERCISES [D].

1. Find the eccentricity of each of the following ellipses :  
 (i.)  $3x^2 + 4y^2 = c^2$ .      (ii.)  $3x^2 + 5y^2 = n^2$ .  
 (iii.)  $25x^2 + 169y^2 = 780^2$ .
2. What is the eccentricity of the ellipse, when A'B is parallel to CF, in fig. of art. 34 ?
3. The focal distances of a point P on an ellipse, whose eccentricity is  $\frac{4}{5}$ , are SP=4, HP=6. Find the coordinates of P referred to the centre as origin.
4. Find the equation to the normal at F, an extremity of the latus rectum.
5. The coordinates of a point P, on an ellipse whose eccentricity is  $\frac{4}{5}\sqrt{2}$ , are  $x' = \frac{27}{8}\sqrt{2}$  and  $y' = \frac{7}{8}\sqrt{238}$ . Find SP and HP.
6. Shew that the equation to the tangent to the ellipse  $2x^2 + 3y^2 = 18$ , inclined to the axis of  $x$  at an angle of  $30^\circ$ , is  $3y = x\sqrt{3} \pm 9$ .
7. Shew that, in an ellipse, if  $a^2 = 2b^2$ , the angle SBH is a right angle.
8. Shew that the lengths of the perpendiculars from the foci of an ellipse, on a tangent inclined to the axis major at an angle  $\phi$ , are  $= a \{e \sin \phi \pm (1 - e^2 \cos^2 \phi)^{\frac{1}{2}}\}$ .
9. Assuming  $\pi a^2$  as the area of the circle described on the major axis of the ellipse, shew that the radius of a circle is a mean proportional between the semi-axes of an ellipse equal in area to the circle.
10. Find the equation to the tangent at F, the extremity of the latus rectum of the ellipse  $\frac{x^2}{9a^2} + \frac{y^2}{4a^2} = 1$ , and the lengths of the intercepts of the tangent on the axes.
11. Two given circles touch each other internally : shew that the locus of the centre of a circle in the space between their circumferences, and touching both, is an ellipse having their centres for its foci.
12. Shew that if  $y$  be the ordinate of any point P on an ellipse, the tangent of the angle A'PA is  $= -\frac{2b^2}{ae^2y}$ .

## THE HYPERBOLA.

53. The hyperbola is the locus of a point, whose distance from a given fixed point is always greater, in a constant ratio, than its distance from a given fixed straight line.

Let  $H$  be the given point, and  $KK'$  the given straight line. Draw  $HD$  perpendicular to  $KK'$ . Let  $P$  be a point on the locus; join  $HP$ , and draw  $PM$  parallel to  $KK'$ , and  $PN$  parallel to  $DH$ .



Take  $D$ , in the line  $KK'$ , as the origin,  $DH$  as the axis of  $x$ ,  $KK'$  as the axis of  $y$ .

Divide  $DH$  at  $A$ , so that  $\frac{AH}{AD} = \frac{HP}{PN}$  = the constant ratio, a value greater than unity, which we may call  $e$ . Let  $DH$  be called  $p$ ; and let  $x, y$  be the coordinates of  $P$ .

We have  $HP = e \cdot PN$ ;  $\therefore HP^2 = e^2 PN^2$ ,

$$\text{or, } PM^2 + HM^2 = e^2 DM^2,$$

$$\text{that is, } y^2 + (x-p)^2 = e^2 x^2;$$

which is the equation to the hyperbola referred to the origin and axes assumed.

54. For the point where the hyperbola meets the axis of  $x$ , we have  $y=0$ , and the equation becomes

$$(x-p)^2 = e^2 x^2;$$

this gives  $x-p = \pm ex$ , or  $(1 \mp e)x = p$ , or  $x = \frac{p}{1 \mp e}$ , signi-

fying that the hyperbola meets the axis at two points, A and A'; but as  $e$  exceeds unity,  $1-e$  is a negative quantity; so that we have DA', measured to the left of D,  $= \frac{p}{e-1}$ , and  $DA = \frac{p}{1+e}$ .

C, the middle point of A'A, is called the centre, and the extremities A, A' are called the vertices, of the hyperbola. Also, the fixed point H is called the focus, and the fixed line KK' the directrix of the hyperbola.

**55.** Let the line A'A  $= 2a$ ; then CA  $= a$ .

Since A' is a point on the locus, we have

$$e = \frac{HA'}{DA'} = \frac{2a + AH}{2a - DA} = \frac{2a + e \cdot DA}{2a - DA};$$

whence are derived the following values:

$$(i.) \quad DA = \frac{a}{e}(e-1);$$

$$(ii.) \quad AH = e \cdot DA = a(e-1);$$

$$(iii.) \quad CD = a - DA = \frac{a}{e};$$

$$(iv.) \quad CH = a + AH = ae;$$

$$(v.) \quad DH = p = CH - CD = ae - \frac{a}{e} = \frac{a}{e}(e^2 - 1).$$

**56.** Suppose the origin to be at A.

In this case, AM takes the place of DM; and if AM be called  $x'$ , then DM  $= DA + x'$ , that is,  $x = x' + a - \frac{a}{e}$ ; whence,

by substitution in the equation

$$y^2 + (x-p)^2 = e^2 x^2,$$

$$\text{we have } y^2 + \left(x' + a - \frac{a}{e} - ae + \frac{a}{e}\right)^2 = \{ex' + a(e-1)\}^2,$$

$$\text{or, } y^2 + \{x' - a(e-1)\}^2 = \{ex' + a(e-1)\}^2,$$

$$\text{or, } y^2 + x'^2 - 2a(e-1)x' = e^2 x'^2 + 2ae(e-1)x',$$

$$\text{or, } y^2 + x'^2 - 2a(e+1)(e-1)x' = e^2 x'^2,$$

$$\therefore y^2 - x'^2(e^2 - 1) = 2ax'(e^2 - 1),$$

$$\text{that is, } y^2 = (e^2 - 1)(2ax' + x'^2).$$

Now, as A is to be accounted the origin, we may remove the accent, and the equation will be

$$y^2 = (e^2 - 1)(2ax + x^2).$$

**57.** Suppose the origin to be at C.

Here, again, we might put  $x'$  for CM, and substitute  $x' - \frac{a}{e}$  for  $x$  in the original equation.

Or, we may at once let  $x$  and  $y$  be the coordinates; then we shall have

$$HM = CM - CH = x - ae;$$

$$DM = CM - CD = x - \frac{a}{e};$$

$$\text{or } e \cdot DM = ex - a;$$

hence, equivalent to  $PM^2 + HM^2 = e^2 DM^2$ , we have

$$y^2 + (x - ae)^2 = (ex - a)^2,$$

$$\text{or, } y^2 = (e^2 - 1)(x^2 - a^2).$$

Now, if we suppose  $x=0$ , we get  $y^2 = a^2(1 - e^2)$ , a negative value, since  $e$  is greater than unity; and therefore, assigning to  $y$  an impossible value, that is, denoting that the curve does not cut the axis of  $y$ . As, however, there will be a point on the curve having for its ordinate  $y' = a\sqrt{e^2 - 1}$ , we may take  $BC = CB' = a\sqrt{e^2 - 1}$ , and call this magnitude  $b$ ;

$$\text{thus } b^2 = a^2(e^2 - 1), \text{ or } e^2 - 1 = \frac{b^2}{a^2}.$$

The equation, therefore, may be written

$$y^2 = \frac{b^2}{a^2}(x^2 - a^2),$$

which is reducible to the more symmetrical form

$$a^2 y^2 - b^2 x^2 = -a^2 b^2, \quad \text{or } \frac{x^2}{a^2} - \frac{y^2}{b^2} = 1;$$

which is the equation to the hyperbola, referred to the centre as origin, and is the one most frequently employed.

**58.** The above equation is convertible into the forms

$$y = \pm \frac{b}{a} \sqrt{x^2 - a^2}, \quad (1)$$

$$x = \pm \frac{a}{b} \sqrt{y^2 + b^2}; \quad (2)$$

which enable us readily to determine the form of the hyperbola.

If  $y=0$ , then (2) gives  $x=\pm a$ ; therefore the hyperbola cuts the axis of  $x$  at the points A, A'.

If  $x$  be supposed a less magnitude than  $+a$  or  $-a$ , then in (1) we shall have  $x^2 - a^2$  negative, and the values of  $y$  are imaginary. Therefore, no point of the hyperbola is situated between A and A'.

If  $x$  be supposed a greater magnitude than  $+a$  or  $-a$ , then (1) shews that for every value of  $x$  there are two values of  $y$ , numerically equal, but contrary in sign. This implies that, for every point above the axis of  $x$ , there is a corresponding point at the same distance below the axis of  $x$ , so that the axis of  $x$  divides the curve symmetrically.

Moreover, as  $x$  increases, the values of  $y$  increase; and when  $x$  becomes indefinitely great, so also does  $y$ .

Accordingly, the hyperbola consists of two opposite and similar branches, respectively extending to the right of A and to the left of A' indefinitely.

It may be here stated that the term *axes*, in reference to the hyperbola, is generally used to denote the lines AA' and BB', and that the former is called the *transverse axis*, the latter the *conjugate axis*.

It may be also mentioned that the ratio denoted by the symbol  $e$  is called the *eccentricity* of the hyperbola.

**59.** The hyperbola being symmetrically situated with respect to the axes AA', BB', there must be two foci, each having its corresponding directrix; and if A'S=AH, and A'E=DA, the curve can be described by means of S and LL', as focus and directrix. (See fig. in 60.)

**60.** The readiest mechanical method of describing an hyperbola is furnished by the following property :


*The difference of the focal distances of any point on the hyperbola is equal to the transverse axis.*

$$\text{For } SP = e \cdot PN' = e(CM + CE) = e\left(x' + \frac{a}{e}\right) = ex' + a,$$

$$HP = e \cdot PN = e(CM - CD) = e\left(x' - \frac{a}{e}\right) = ex' - a,$$

$$\therefore SP - PH = 2a.$$

Accordingly, an hyperbola may be constructed as follows :



Let a ruler SR revolve round S in the plane of the paper, and let a string HPR, fastened to R and H, be of such length, that  $SR - HPR = 2a$ ; then with a pencil, P, keep the string stretched against SR as the ruler revolves, and the point will trace out a portion of the hyperbola of which S and H are the foci, since  $SP - PH$  will be constantly  $= 2a$ .

**61.** The equation  $y^2 = \frac{b^2}{a^2}(x^2 - a^2)$  may be put in the form

$$y^2 = \frac{b^2}{a^2}(x+a)(x-a);$$

hence, if PM be any ordinate, and C the origin,

$$PM^2 = \frac{BC^2}{CA^2} \cdot A'M \cdot MA,$$

$$\text{or } PM^2 : A'M \cdot MA :: BC^2 : CA^2.$$

62. The double ordinate FHF', passing through the focus, is called the *latus rectum*. To find its length, we have  $x = CH = ae$ , which substituted for  $x$  in the equation

$$y^2 = \frac{b^2}{a^2} (x^2 - a^2), \text{ gives}$$

$$y^2 = \frac{b^2}{a^2} (a^2 e^2 - a^2) = b^2 (e^2 - 1) = b^2 \cdot \frac{b^2}{a^2} = \frac{b^4}{a^2};$$

$$\text{hence } y = FH = \frac{b^2}{a}, \text{ and } FHF' = \frac{2b^2}{a}.$$

#### TANGENT AND NORMAL TO AN HYPERBOLA.

63. If we compare the equations to the ellipse and hyperbola,

$$\text{that for the ellipse being } y^2 = \frac{b^2}{a^2} (a^2 - x^2),$$

$$\begin{aligned} \text{that for the hyperbola } y^2 &= \frac{b^2}{a^2} (x^2 - a^2) \\ &= -\frac{b^2}{a^2} (a^2 - x^2), \end{aligned}$$

it will appear that the latter may be derived from the former by writing  $-b^2$  for  $b^2$ . Hence

To find the equation to the tangent at a given point  $(x', y')$  on an hyperbola :

Instead of repeating the procedure in art. 45, we need only write  $-b^2$  for  $b^2$  in the equation there found, and we have

$$y - y' = \frac{b^2 x'}{a^2 y'} (x - x'),$$

$$\text{or } a^2 y y' - b^2 x x' = -a^2 b^2.$$

64. To express the tangent to an hyperbola in terms of the tangent of the angle which the line makes with the axis of  $x$ .

Here the required expression may be at once derived from that in art. 46; thus,

$$y = mx + \sqrt{a^2 m^2 - b^2};$$





**66.** *The focal distances of any point on the hyperbola make equal angles with the tangent at that point.*

This may be proved by the method employed in art. 48. And hence also the angle between the focal distances of any point on the hyperbola is bisected by the normal at that point; or  $SPT = HPT$ .

**67.** *To find the locus of the intersection of a tangent to the hyperbola with the perpendicular drawn to it from either focus.*

By pursuing the method employed in art. 50, the required locus will be ascertained to be a circle of which the transverse axis is a diameter; and the result will, as in that article, suggest a geometrical method of drawing a tangent to an hyperbola from any external point.

**68.** *To determine the perpendicular,  $p$ , from the centre of an hyperbola as origin, to the tangent at any point on the curve, in terms of the angle  $\theta$ , which  $p$  makes with the axis of  $x$ .*

In the expression found in art. 51, substitute  $-b^2$  for  $b^2$ , and the equation determining  $p$  will be

$$\begin{aligned} p^2 &= a^2 \cos^2 \theta - b^2 \sin^2 \theta, \\ \text{or } p^2 &= a^2 \cos^2 \theta - a^2(e^2 - 1) \sin^2 \theta, \\ &= a^2(\cos^2 \theta + \sin^2 \theta) - a^2 e^2 \sin^2 \theta, \\ \text{that is, } p^2 &= a^2(1 - e^2 \sin^2 \theta); \end{aligned}$$

which is the same result as that for the ellipse.

**69.** All the properties of the hyperbola shewn in the preceding articles on that curve have been found similar to those of the ellipse. There are peculiar properties of the hyperbola, of which, in this introductory treatise, we do not consider the investigation to be necessary. We will, however, direct the attention of the student to one remarkable property of the hyperbola,—that of the asymptotes.

The equation to the hyperbola, viz.,

$$y^2 = \frac{b^2}{a^2} (x^2 - a^2),$$

may be put in the form


$$y^2 = \frac{b^2 x^2}{a^2} \left(1 - \frac{a^2}{x^2}\right);$$

$$\text{hence } y = \pm \frac{bx}{a} \sqrt{\left(1 - \frac{a^2}{x^2}\right)}.$$

Now, as  $x$  increases, the value of  $\frac{a^2}{x^2}$  diminishes, so that that value becomes zero when, but not until,  $x$  becomes infinite; and then

$$y = \pm \frac{b}{a} x.$$

If, therefore, through the centre  $C$ , as origin, two straight lines,  $CL$ ,  $CK$ , be drawn, making with the axis of  $x$  angles  $LCX$ ,  $KCX$ , whose tangents are, respectively,  $\frac{b}{a}$  and  $-\frac{b}{a}$ , these lines will continually approach the curve, and yet never meet it. For the equations to the lines are



$$y = \frac{b}{a} x, \text{ and } y = -\frac{b}{a} x,$$

and we have seen that  $y$  cannot become  $= \pm \frac{b}{a} x$  till  $x$  becomes infinite.

The lines  $CL$ ,  $CK$  are called *Asymptotes* to the hyperbola: the word asymptote, of Greek origin, signifying *without coincidence* or contact.

**70.** When the angle  $LCX$  is half a right angle, the tangent  $\frac{b}{a} = 1$ , or  $b = a$ ; and the value  $b^2 = a^2(e^2 - 1)$  gives  $e^2 = 2$ . The curve is then called an *equilateral* hyperbola, because

of the equality of the axes, and also a *rectangular* hyperbola, because of LCL', the angle between the asymptotes, being a right angle.

**71.** To find the equation to the rectangular hyperbola referred to the asymptotes as axes.

In the equation to the hyperbola,

$$a^2y^2 - b^2x^2 = -a^2b^2,$$

let  $b=a$ ; then

$$x^2 - y^2 = a^2.$$

Now, let P be a point on the rectangular hyperbola, CX, CY the original axes, CX<sub>1</sub>, CY<sub>1</sub> the axes coincident with the asymptotes.

Let CM =  $x$ , PM =  $y$ , CM' = RM' =  $x'$ , PM' =  $y'$ .

$$\begin{aligned} x &= CM = CN + NM \\ &= CM' \frac{CN}{CM'} + P'N \frac{NM}{P'N} \\ &= x' \cos 45^\circ + y' \cos 45^\circ \\ &= \frac{1}{2}\sqrt{2}(x' + y'); \end{aligned}$$

$$y = PM = PR \frac{PM}{PR} = (y' - x') \sin 45^\circ = \frac{1}{2}\sqrt{2}(y' - x');$$


$$\therefore x^2 - y^2 = \frac{1}{2}(x' + y')^2 - \frac{1}{2}(y' - x')^2 = a^2,$$

$$\text{that is, } 4x'y' = 2a^2,$$

or, suppressing the accents,

$$xy = \frac{1}{2}a^2,$$

the equation required.



**72.** We will now propose a few Exercises on the hyperbola without an introduction of worked Examples.

## EXERCISES [E].

1. The eccentricity of an hyperbola is 2 : Shew that the equation to the curve is  $y=3(x^2-a^2)$ .

2. Required the eccentricity and latus rectum of an hyperbola whose equation is  $2y^2=3(x^2+2n^2)$ .

3. Shew that the tangent at A, the vertex of an ellipse or of an hyperbola, is at right angles to the transverse axis.

4. The tangent to an hyperbola at P cuts the transverse axis at T, and the tangent at the vertex A intercepts CP at E : Shew that the straight line TE is parallel to the straight line AP.

5. What relation must subsist between the constants, in order that the line  $\frac{y}{m} - \frac{x}{n} = 1$  may be a tangent to the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  ?

6. If an ellipse and an hyperbola have the same foci, the locus of the intersection of tangents to them, at right angles to each other, is a circle.

7. Find the locus of the intersection of tangents drawn from the fixed points A and B to a circle touching the straight line AB at a fixed point.

8. If P be any point on an hyperbola, shew that

$$\tan \frac{1}{2} \text{PSH} \tan \frac{1}{2} \text{PHS} = \frac{e-1}{e+1}.$$

9. Through the extremities of PP', a double ordinate of an ellipse, are drawn the straight lines A'PQ, P'AQ, meeting at Q : Shew that the locus of Q is an hyperbola having the same axes as the ellipse.

10. A circle is described, having the focus of an hyperbola as centre, and a diameter equal to the conjugate axis : Shew that the asymptotes are tangents to the circle at the points where they intersect the nearer directrix.

## THE PARABOLA.

**73.** The parabola is the locus of a point whose distance from a given fixed point is always equal to its distance from a given fixed straight line.

Let  $S$  be the given point, and  $KK'$  the given line. Draw  $SD$  perpendicular to  $KK'$ . Let  $P$  be a point on the locus; join  $SP$ , and draw  $PM$  parallel to  $KK'$ , and  $PN$  parallel to  $DS$ .

Take  $D$  as the origin,  $DS$  as the axis of  $x$ ,  $KK'$  as the axis of  $y$ .


Bisect  $DS$  at  $A$ ; suppose  $AS=a$ ; and let  $x, y$  be the coordinates of  $P$ .

We have  $SP=PN$ ;  $\therefore SP^2=PN^2$ ;  
or  $PM^2+SM^2=DM^2$ ;

$$\text{that is, } y^2 + (x-2a)^2 = x^2,$$

$$\text{or, } y^2 = 4a(x-a);$$

which is the equation to the parabola, referred to the origin and axes assumed.



**74.** For the point where the parabola meets the axis of  $x$ , we have  $y=0$ , and the equation becomes  $x=a$ , signifying that the curve meets the axis at one point  $A$ , so that  $DA=AS$ .

$A$  is called the vertex,  $AX$  the axis,  $S$  the focus, and  $KK'$  the directrix, of the parabola.

**75.** Suppose the origin to be at  $A$ .

In this case,  $AM$  takes the place of  $DM$ ; and if  $AM$  be called  $x'$ , then  $DM=x'+a$ , that is,  $x=x'+a$ ; whence, by substitution in the equation  $y^2=4a(x-a)$ , we have

$$y^2=4ax'.$$

Now, as  $A$  is to be accounted the origin, we may remove the accent, and we shall have

$$y^2=4ax,$$

which is the equation to the parabola referred to the vertex as origin, and is the one most frequently employed.

**76.** The above equation is convertible into the form

$$y = \pm 2\sqrt{ax},$$

which enables us readily to determine the form of the parabola.

If  $x$  be supposed negative, the values of  $y$  are imaginary. Therefore no point on the parabola is situated on the left of the origin.

For every positive value of  $x$  there are two values of  $y$ , numerically equal, but contrary in sign. This implies that for every point above the axis of  $x$  there is a corresponding point at the same distance below the axis of  $x$ ; so that the axis of  $x$  divides the curve symmetrically.


Moreover, as  $x$  increases, the values of  $y$  increase, and when  $x$  becomes indefinitely great, so also does  $y$ .

Accordingly, the parabola is a non-central curve, extending indefinitely to the right of A.

**77.** The focal distance of any point P on the parabola, being equal to its distance from the directrix, we have

$$SP = AM + AS, \text{ or } SP = x + a.$$

The double ordinate LSL' passing through the focus is called the *latus rectum*. To find its length, we have  $x = AS = a$ ;  $y^2 = LS^2 = 4ax = 4a^2$ ; hence  $y = \pm 2a$ ;  $\therefore LSL' = 4a$ .



#### TANGENT AND NORMAL TO A PARABOLA.

**78.** The tangent to a parabola may be regarded as a secant passing through two coincident points on the curve. (See art. 27.)

*To find the equation to the tangent at a given point on a parabola.*

Let  $x', y'$  be the coordinates of P the given point,  $x'', y''$  those of another point P' on the curve, near the given point.

The equation to the straight line PP' is

$$y - y' = \frac{y'' - y'}{x'' - x'} (x - x'). \quad (1)$$

The equation to the parabola referred to the vertex as origin is

$$y^2 = 4ax.$$

Accordingly, since  $(x', y')$  and  $(x'', y'')$  are points on the curve,

$$y'^2 = 4ax', \text{ and } y''^2 = 4ax'';$$

$$\therefore y''^2 - y'^2 = 4a(x'' - x'),$$

$$\text{or } (y'' + y')(y'' - y') = 4a(x'' - x');$$

$$\therefore \frac{y'' - y'}{x'' - x'} = \frac{4a}{y'' + y'};$$

hence the 1st equation, by substitution, will become

$$y - y' = \frac{4a}{y'' + y'} (x - x'). \quad (2)$$

Now, when the point  $(x'', y'')$  coincides with the point  $(x', y')$ , we have  $x'' = x'$ , and  $y'' = y'$ ; in which case (2)

becomes 
$$y - y' = \frac{4a}{2y'} (x - x'),$$

$$\text{that is, } y - y' = \frac{2a}{y'} (x - x'),$$

$$\text{or } yy' - 2ax + 2ax' = y'^2 = 4ax';$$

$$\therefore yy' = 2a(x + x').$$

**79.** *To express the tangent to a parabola in terms of the tangent of the angle which the line makes with the axis of  $x$ .*

We have seen that the tangent of the angle which the line makes with the axis of  $x$  is  $\frac{2a}{y'}$ .



From the equation  $yy' = 2a(x + x')$

$$\text{we have } y = \frac{2a}{y'}x + \frac{2ax'}{y'},$$

$$\text{or } y = \frac{2a}{y'}x + \frac{4ax'}{2y'},$$

$$\text{that is, } y = \frac{2a}{y'}x + \frac{y'^2}{2y'},$$

$$\text{or } y = \frac{2a}{y'}x + \frac{y'}{2}.$$

Let  $\frac{2a}{y'} = m$ ;  $\therefore \frac{y'}{2a} = \frac{1}{m}$ , or  $\frac{y'}{2} = \frac{a}{m}$ ; hence by substitution

$$y = mx + \frac{a}{m};$$

which is the equation to the tangent in terms of the tangent of the angle which the line makes with the axis of the parabola.

**80.** *To find the equation to the normal at any point on a parabola.*

Let  $(x', y')$  be the point of contact of a tangent to the parabola.

The equation to the tangent is

$$y = \frac{2a}{y'}x + \frac{y'}{2}.$$

The equation to the normal, as a straight line through  $(x', y')$ , is of the form

$$y - y' = m'(x - x');$$

and the condition of its being perpendicular to the tangent requires that  $m' = -\frac{1}{m} = -\frac{y'}{2a}$ ; hence the equation to the normal at  $(x', y')$  is

$$y - y' = -\frac{y'}{2a}(x - x').$$

At G, where the normal PG meets the axis,  $y=0$ , and the above equation gives


$$-y' = -\frac{y'}{2a} (x-x'),$$

$$\text{or } 1 = \frac{1}{2a} (x-x'),$$

$$\text{or } x-x' = 2a,$$

that is,  $AG - AM = MG = 2a$ .

At T, where the tangent TP meets the axis,  $y=0$ , and the equation to the tangent, viz.,  $yy' = 2a(x+x')$ , becomes  $x = -x'$ , that is,  $AM = AT$ .



Hence also  $ST = x + a = SP$ .

Therefore the axis of a parabola, and the focal distance of any point on the curve to which a tangent is drawn, make equal angles with the tangent, and also equal angles with the normal; consequently  $SG = SP$ .

**81.** To find the locus of the intersection of a tangent to the parabola with the perpendicular drawn to it from the focus.

The equation to TP is

$$y = mx + \frac{a}{m}. \quad (1)$$

The equation to SR drawn from the focus whose coordinates are  $a, 0$ , is of the form

$$y = m'(x-a),$$

and that this line may be perpendicular to the former we put  $m' = -\frac{1}{m}$ , or

$$y = -\frac{1}{m} (x-a); \quad (2)$$

$$\text{hence } mx + \frac{a}{m} = -\frac{1}{m} (x-a),$$

$$\text{or } m^2x + a = a - x,$$

$$\text{or } (1 + m^2)x = 0;$$

but  $1 + m^2$  is not  $= 0$ ;  $\therefore x = 0$ ;

or, the locus of the intersection of a tangent to the parabola with the perpendicular on it from the focus is the axis of  $y$ .

**82.** We shall here propose a few Exercises on the parabola.

### EXERCISES [F].

1. A straight line AL is drawn from the vertex of a parabola to an extremity of the latus rectum: Find the angle which AL makes with a tangent at L.

2. Find the length of TP, a tangent to a parabola at the point P, inclined to the axis at an angle  $PTX = 60^\circ$ .

3. LL' being the latus rectum of a parabola whose vertex is A: Find the diameter of a circle described about the triangle LAL'.

4. If the tangent at a point P on a parabola be inclined to the axis at an angle of  $30^\circ$ , shew that the focal distance is equal to the latus rectum.

5. A tangent is drawn to a parabola at L, an extremity of the latus rectum: Find the length of the normal chord LGQ'.

6. AB is a fixed diameter of a circle whose centre is C; and DE is any chord parallel to AB: Find the locus of the intersection of CD with a straight line AQ drawn to the middle point of DE.

7. PSP' is any focal chord of a parabola, the straight line AP cutting the latus rectum at Q: Shew that the straight line QP' is bisected by the axis.

8. LS is the semi-latus rectum of a parabola APL, whose vertex is A and focus S; and D is the point where the axis meets the directrix; a straight line from D through P

meets the semi-latus rectum at  $N$ , and  $NQ$  is drawn parallel to  $DS$ , and  $SPQ$  is drawn to meet  $NQ$ : Shew that the locus of  $Q$  is a circle.


9. Two tangents are drawn to a parabola, and are inclined to the axis at angles the difference of whose cotangents is equal to a constant quantity  $d$ : Shew that the locus of the intersection of the tangents is a parabola equal to the original parabola.

## TRANSFORMATION OF COORDINATES.

**83.** It is often convenient to alter given coordinates with respect to origin, or direction, or both. This can be done with either oblique or rectangular axes. We confine our illustrations to the latter.

*To transfer the origin of coordinates to another point, without changing the direction of the axes.*

Let  $O'X'$ ,  $O'Y'$  be new axes parallel to the original axes  $OX$ ,  $OY$ . Let the coordinates of the new origin referred to the old origin be  $OA=h$ ,  $O'A=k$ . Let  $x$ ,  $y$  be the original coordinates of a point  $P$ , and  $x'$ ,  $y'$  the new coordinates of the same point.



$$x = OM = AM + OA = x' + h;$$


$$y = PM = PM' + M'M = y' + k;$$

which are the values of  $x$  and  $y$  that must be substituted in order to express what the equation to any locus becomes when referred to the new axes.

**84.** *To change one system of rectangular axes to another with the same origin.*

Let  $OX'$ ,  $OY'$  be new rectangular axes, having the same origin as  $OX$ ,  $OY$ , the original axes. Let  $x$ ,  $y$  be the original coordinates of a point  $P$ , and  $x'$ ,  $y'$  the new coordinates of the point.

Draw  $PM$ ,  $PM'$  parallel to  $OY$ ,  $OY'$ , and  $M'N$ ,  $M'R$  parallel to  $OY$ ,  $OX$ .



The axes have been turned round  $O$ , through an angle  $XOX'$ ; let that angle  $= \theta$ ;  
then

$$x = OM = ON - RM',$$

$$y = PM = PR + M'N;$$

but the angle  $RPM' = XOX' = \theta$ ;

$$\therefore RM' = PM' \sin \theta; \quad \text{and} \quad PR = PM' \cos \theta;$$

$$ON = OM' \cos \theta; \quad \text{and} \quad M'N = OM' \sin \theta.$$

$$\therefore x = x' \cos \theta - y' \sin \theta;$$

$$y = x' \sin \theta + y' \cos \theta;$$

which are the values of  $x$  and  $y$  that must be substituted in order to express what the equation to any locus becomes when referred to the new axes.

To make this case include a transfer of the origin to the point  $(h, k)$ , we have only to substitute  $x'' + h$  for  $x'$ , and  $y'' + k$  for  $y'$ .

**85.** The general equation of the second degree is

$$Ax^2 + Bxy + Cy^2 + Dx + Ey + F = 0;$$

and it most frequently represents some curve.

It may, however, represent two straight lines. Thus, if we have

$$6x^2 - 11xy + 3y^2 - 3x + 8y - 3 = 0,$$

then solving this as a quadratic for  $y$ , we shall obtain as the two values of  $y$ ,

$$y = 3x - 3, \text{ and } = \frac{2}{3}x + \frac{1}{3},$$

shewing that the given equation is the product of

$$y - 3x + 3 = 0$$

$$\text{and } 3y - 2x - 1 = 0,$$

and therefore represents two straight lines.

We proceed to shew how transformation of coordinates may be applied for the reduction of the general equation of the second degree to the forms by which loci are readily distinguished.

**86.** Let us begin by examining how the terms involving the simple powers of  $x$  and  $y$  may be removed.

If we transfer the origin to a point  $(h, k)$ , without altering the direction of the rectangular axes; that is, if we substitute  $x' + h$  for  $x$ , and  $y' + k$  for  $y$ , we shall have

$$\begin{aligned} & Ax'^2 + 2Ahx' + Ah^2 + Bx'y' + Bkx' + Bhy' + Bhk \\ & + Cy'^2 + 2Cky' + Ck^2 + Dx' + Dh + Ey' + Ek + F = 0; \end{aligned}$$

that is,

$$\begin{aligned} & Ax'^2 + Bx'y' + Cy'^2 + (2Ah + Bk + D)x' + (Bh + 2Ck + E)y' \\ & + Ah^2 + Bhk + Ck^2 + Dh + Ek + F = 0. \end{aligned}$$

Now, to try to get rid of the terms of  $x'$  and  $y'$ , let us equate their coefficients to zero; thus,

$$2Ah + Bk + D = 0; \quad \text{and } Bh + 2Ck + E = 0;$$

$$\therefore k = \frac{-2Ah - D}{B} = \frac{-Bh - E}{2C}; \quad (1)$$

$$h = \frac{-Bk - D}{2A} = \frac{-2Ck - E}{B}. \quad (2)$$

$$\text{From (1) we have } h = \frac{2C \cdot D - B \cdot E}{B^2 - 4A \cdot C}.$$

$$\text{From (2) we have } k = \frac{2A \cdot E - B \cdot D}{B^2 - 4A \cdot C}.$$

It appears then that such values of  $h$  and  $k$  can be assigned as will make the coefficients of  $x$  and  $y = 0$ , except when  $B^2 - 4A \cdot C = 0$ .

87. When  $B^2 - 4A \cdot C = 0$ , or  $B^2 = 4A \cdot C$ , we do not get rid of the simple powers of  $x$  and  $y$  by equating their coefficients to zero as above; but we can get rid of the product of the variables by putting

$$x = x' \cos \theta - y' \sin \theta,$$

$$y = x' \sin \theta + y' \cos \theta.$$

The general equation then becomes

$$\begin{aligned} A(x'^2 \cos^2 \theta - 2x'y' \cos \theta \sin \theta + y'^2 \sin^2 \theta) + B(x'^2 \sin \theta \cos \theta - \\ x'y' \sin^2 \theta + x'y' \cos^2 \theta - y'^2 \sin \theta \cos \theta) + C(x'^2 \sin^2 \theta + \\ 2x'y' \sin \theta \cos \theta + y'^2 \cos^2 \theta) + D(x' \cos \theta - y' \sin \theta) + \\ E(x' \sin \theta + y' \cos \theta) + F = 0; \end{aligned}$$

or, since  $2 \sin \theta \cos \theta = \sin 2\theta$ , and  $\cos^2 \theta - \sin^2 \theta = \cos 2\theta$ ,

$$\left. \begin{aligned} &x'^2(A \cos^2 \theta + \frac{1}{2}B \sin 2\theta + C \sin^2 \theta) \\ &+ x'y' \{(C - A) \sin 2\theta + B \cos 2\theta\} \\ &+ y'^2(A \sin^2 \theta - \frac{1}{2}B \sin 2\theta + C \cos^2 \theta) \\ &+ x'(D \cos \theta + E \sin \theta) \\ &+ y'(E \cos \theta - D \sin \theta) \\ &+ F \end{aligned} \right\} = 0.$$

Now, the tangent of an angle may be of any magnitude; and therefore we may put

$$\frac{B}{A - C} = \tan 2\theta = \frac{\sin 2\theta}{\cos 2\theta},$$

whereby we make  $-B \cos 2\theta = (C - A) \sin 2\theta$ ,

and thus, by substitution in the coefficient of  $x'y'$ , that coefficient will become zero.

We can, therefore, always get rid of the product of the variables by turning the axes through an angle  $\theta$  such that

$$\tan 2\theta = \frac{B}{A - C}.$$

Let us now seek the corresponding values of the coefficients of  $x'^2$  and  $y'^2$ .

$$2A \cos^2 \theta + 2C \sin^2 \theta = A(2 - 2 \sin^2 \theta) + 2C \sin^2 \theta;$$

but, by Trigonometry,  $2 \sin^2 \theta = 1 - \cos 2\theta$ ; thus making the expression become

$$\begin{aligned} &= A(1 + \cos 2\theta) + C(1 - \cos 2\theta) \\ &= A + C + (A - C) \cos 2\theta; \\ \therefore A \cos^2 \theta + C \sin^2 \theta &= \frac{1}{2} \{A + C + (A - C) \cos 2\theta\}. \end{aligned}$$

Similarly,

$$A \sin^2 \theta + C \cos^2 \theta = \frac{1}{2} \{A + C - (A - C) \cos 2\theta\}.$$

Hence the terms involving  $x'^2$  and  $y'^2$  become

$$\begin{aligned} &x'^2 \cdot \frac{1}{2} \{A + C + (A - C) \cos 2\theta + B \sin 2\theta\} \\ &+ y'^2 \cdot \frac{1}{2} \{A + C - (A - C) \cos 2\theta - B \sin 2\theta\}. \end{aligned}$$

$$\text{Now, } \tan^2 2\theta = \frac{\sin^2 2\theta}{\cos^2 2\theta} = \frac{B^2}{(A - C)^2};$$

$$\therefore B \sin 2\theta = \frac{B^2}{\sqrt{(A - C)^2 + B^2}},$$

$$(A - C) \cos 2\theta = \frac{(A - C)^2}{\sqrt{(A - C)^2 + B^2}};$$

so that the terms involving  $x'^2$  and  $y'^2$  become

$$\begin{aligned} &x'^2 \cdot \frac{1}{2} \{A + C + \sqrt{(A - C)^2 + B^2}\} \\ &+ y'^2 \cdot \frac{1}{2} \{A + C - \sqrt{(A - C)^2 + B^2}\}; \end{aligned}$$

where it will be observed that the product of the coefficients is  $\frac{1}{4}(4A \cdot C - B^2)$ ; if, therefore,  $4A \cdot C - B^2 = 0$ , one of these coefficients must vanish.

In this case, if the coefficient of  $x'^2 = 0$ , then the general equation, having lost  $x'^2$  and  $x'y'$ , becomes

$$C'y'^2 + D'x' + E'y' + F = 0,$$

which may be written

$$y'^2 + \frac{E'}{C'} y' + \frac{D'}{C'} x' + \frac{F}{C'} = 0,$$

$$\text{or, } \left(y' + \frac{E'}{2C'}\right)^2 = -\frac{D'}{C'} x' - \frac{F}{C'} + \frac{E'^2}{4C'},$$

$$\text{or, } \left(y' + \frac{E'}{2C'}\right)^2 = -\frac{D'}{C'} \left(x' + \frac{F}{D'} - \frac{E'^2}{4C'D'}\right).$$



Let  $-\frac{E'}{2C'}$  be called P, and let  $\frac{E'^2}{4C'D'} - \frac{F'}{D'}$  be called R;

then,  $(y' - P)^2 = -\frac{D'}{C'} (x' - R)$ ;

hence, if D be not = 0, and the origin be transferred to the point (R, P),

writing  $x'' + R$  for  $x'$ , and  $y'' + P$  for  $y'$ , we shall have

$$y''^2 = -\frac{D'}{C'} x'',$$

which is the equation to a parabola, the coordinates to the vertex being R and P.

**88.** We will now proceed to Examples and Exercises on the use of those expedients that we have found available for the transformation of coordinates.

#### EXAMPLES.

*Ex.* (1). Transform the equation  $x + 2\sqrt{xy} + y = c$ , by turning the axes of the coordinates through half a right angle.

Here  $\theta = 45^\circ$ , and  $\sin \theta = \cos \theta = \frac{1}{2}\sqrt{2}$ .

Putting  $x = x' \cos \theta - y' \sin \theta$ , and  $y = x' \sin \theta + y' \cos \theta$ , we have  $x = \frac{1}{2}\sqrt{2}(x' - y')$ ; and  $y = \frac{1}{2}\sqrt{2}(x' + y')$ ;

$\therefore x + y + 2\sqrt{xy} = c$  becomes

$$\begin{aligned}\sqrt{2}x' + 2\sqrt{\frac{1}{2}(x'^2 - y'^2)} &= c; \\ \text{or, } (\sqrt{2}x' - c)^2 &= 2x'^2 - 2y'^2; \\ \text{or, } y'^2 &= c\sqrt{2}x' - \frac{1}{2}c^2.\end{aligned}$$

*Ex.* (2). Transform the equation

$$5x^2 - 4xy + 2y^2 - 3x = 0,$$

by removing the simple power of  $x$  and the product of the variables.

Here  $A=5$ ,  $B=-4$ ,  $C=2$ ,  $D=-3$ ,  $E=0$ ,  $F=0$ ; and  $B^2 - 4A \cdot C$  being  $=16 - 40 = -24$ , we can get rid of the

simple power of  $x$  by transferring the origin to the point  $(h, k)$ , taking

$$h = \frac{2C \cdot D - B \cdot E}{B^2 - 4A \cdot C} = \frac{-12}{-24} = \frac{1}{2},$$

$$k = \frac{2A \cdot E - B \cdot D}{B^2 - 4A \cdot C} = \frac{-12}{-24} = \frac{1}{2}.$$

Thus for  $x$  take  $x' + \frac{1}{2}$ , and for  $y$  take  $y' + \frac{1}{2}$ ; then,

$$5(x' + \frac{1}{2})^2 - 4(x' + \frac{1}{2})(y' + \frac{1}{2}) + 2(y' + \frac{1}{2})^2 - 3(x' + \frac{1}{2}) = 0;$$

$$\text{or } 20x'^2 - 16x'y' + 8y'^2 - 3 = 0.$$

Now, to remove the product of the variables, take  $\tan 2\theta$

$$= \frac{B}{A - C} = \frac{-4}{5 - 2} = -\frac{4}{3}.$$

$$\text{By Trigonometry, } \tan \theta = \frac{\sqrt{(1 + \tan^2 2\theta)} - 1}{\tan 2\theta} = \frac{-5 - 3}{-4} = 2;$$

$$\therefore \sin \theta = \frac{2}{5}\sqrt{5}, \text{ and } \cos \theta = \frac{1}{5}\sqrt{5}.$$

Hence, putting  $x'' \cos \theta - y'' \sin \theta$  for  $x'$ ,

$$x'' \sin \theta + y'' \cos \theta \text{ for } y',$$

we have

$$\begin{aligned} 20(\frac{1}{5}\sqrt{5}x'' - \frac{2}{5}\sqrt{5}y'')^2 &= 4x''^2 - 16x''y'' + 16y''^2 \\ 8(\frac{2}{5}\sqrt{5}x'' + \frac{1}{5}\sqrt{5}y'')^2 &= 6\frac{2}{5}x''^2 + 6\frac{2}{5}x''y'' + 1\frac{3}{5}y''^2 \\ -16(\frac{1}{5}\sqrt{5}x'' - \frac{2}{5}\sqrt{5}y'')(\frac{2}{5}\sqrt{5}x'' + \frac{1}{5}\sqrt{5}y'') &= -6\frac{2}{5}x''^2 + 9\frac{2}{5}x''y'' + 6\frac{2}{5}y''^2 \\ \text{sum} &= 4x''^2 + 24y''^2 = 3; \end{aligned}$$

or, suppressing the accents,

$$\frac{x^2}{\frac{3}{4}} + \frac{y^2}{\frac{1}{8}} = 1,$$

which is the equation to an ellipse whose axes are  $2a = \sqrt{3}$ , and  $2b = \frac{1}{2}\sqrt{2}$ .

*Ex. (3).* Required the latus rectum, and the coordinates of the vertex, of the parabola whose equation is

$$x^2 - 4xy + 4y^2 - 4x - 2y + 10 = 0.$$

Here  $A=1$ ,  $B=-4$ ,  $C=4$ ,  $D=-4$ ,  $E=-2$ ,  $F=10$ .

$B^2 - 4AC$  being  $=0$ , we cannot remove the origin. Let us, however, turn the axes through an angle  $\theta$  such that

$$\tan 2\theta = \frac{B}{A - C} = \frac{4}{3}.$$

We have  $\tan \theta = \frac{1}{2}$ ,  $\therefore \sin \theta = \frac{1}{\sqrt{5}}$ , and  $\cos \theta = \frac{2}{\sqrt{5}}$ .

Hence, putting  $\frac{2}{\sqrt{5}}x' - \frac{1}{\sqrt{5}}y'$  for  $x$ ,

$\frac{1}{\sqrt{5}}x' + \frac{2}{\sqrt{5}}y'$  for  $y$ ,

the given equation is reduced to

$$5y'^2 - 2\sqrt{5}x' + 10 = 0,$$

where  $C'=5$ ,  $D'=-2\sqrt{5}$ ,  $E'=0$ ,  $F=10$ ;

hence the formula

$$\left(y' + \frac{E'}{2C'}\right)^2 = -\frac{D'}{C'}\left(x' + \frac{F}{D'} - \frac{E'^2}{4C'D'}\right)$$

gives

$$y'^2 = \frac{2}{\sqrt{5}}(x' - \sqrt{5}),$$

showing the latus rectum  $4a = \frac{2}{\sqrt{5}}$ , and the coordinates of the vertex  $-\sqrt{5}$  and 0.

*Ex. (4).* Trace the curve  $y = x - x^2$ .

We have here  $x^2 + y - x = 0$ . Substitute  $y'$  for  $x$  and  $x'$  for  $y$ ; then

$$y'^2 + x' - y' = 0,$$

where  $C'=1$ ,  $D'=1$ ,  $E'=-1$ ,  $F=0$ ;

hence the formula in the preceding example gives


$$\left(y' - \frac{1}{2}\right)^2 = -\left(x' - \frac{1}{4}\right);$$

now replacing  $y'$  by  $x$  and  $x'$  by  $y$ , we have

$$\left(x - \frac{1}{2}\right)^2 = -\left(y - \frac{1}{4}\right),$$

which is the equation to a parabola, having its axis parallel to that of  $y$ , the coordinates of its vertex  $\frac{1}{2}$  and  $\frac{1}{4}$ , and its latus rectum 1.

*Ex. (5).*  $DQ=l$  is a horizontal line, and  $DO=p$  a perpen-



dicular erected upon it. A cord OPQ, fixed at O, is drawn over the point Q, while at P a weight, freely suspended, slides along the cord. Required the locus of P.

Let the coordinates of Q be  $ON=l$ ,  $QN=p$ ; and let those of P be  $OM=x$ ,  $PM=y$ .

The triangles OPM, EPQ, being similar,  $\frac{EQ}{OM} = \frac{PE}{PM}$ ,

that is,  $\frac{l-x}{x} = \frac{y-p}{y}$ ,

$$\text{or } 2xy - px - ly = 0. \quad (1)$$

Here  $A=0$ ,  $B=2$ ,  $C=0$ ,  $D=-p$ ,  $E=-l$ ; hence  $h=\frac{1}{2}l$ , and  $k=\frac{1}{2}p$ ; and if we put  $x'+\frac{1}{2}l$  for  $x$ , and  $y'+\frac{1}{2}p$  for  $y$ , we have, by substitution in (1),

$$x'y' = \frac{1}{4}lp,$$

which is the equation to an hyperbola whose asymptotes are the axes of  $x$  and  $y$ , and in the present instance rectangular.

Accordingly, since  $\frac{1}{4}lp = \frac{1}{4}ON \cdot QN$ , the locus of P is an equilateral hyperbola whose centre is C the middle point of the diagonal OQ, and its transverse axis A'A making with DQ an angle QFA =  $45^\circ$ .

### EXERCISES [G].

1. Shew that the equation  $3xy - 4x = a^2$  will become  $x^2 - 9y^2 = 2a^2$ , by turning the axes through an angle whose tangent is 3.

2. The equation to a rectangular hyperbola being  $x^2 - y^2 = a^2$ ; transform that equation by turning the axes in the negative direction through half a right angle.

3. Find the locus represented by the equation

$$4x^2 + 12xy + 9y^2 - 1 = 0.$$

4. Find the axes of the curve  $x^2 + xy + y^2 = 1$ .

5. What is represented by the equation

$$9x^2 - 30xy + 25y^2 + 21x - 35y + 10 = 0?$$

6. Determine the axes of the curve which is the locus of the equation

$$x^2 - 6xy + y^2 - 6x + 2y + 5 = 0.$$

7. What is represented by the equation

$$y^2 - 2xy - x^2 + 2 = 0?$$

8. Determine the nature and position of the curve represented by the equation

$$x^2 - 2xy + y^2 - 8x + 16 = 0.$$

9. What is the geometrical construction of the following equation :

$$y^2 - 2xy + x^2 - 2y - 2x - 3 = 0?$$

10. Find the axes and position of the curve represented by the equation

$$3x^2 - 2xy + y^2 - 4x + 2y - 3 = 0.$$

11. The base of a triangle is  $= 2n$ , and the difference of the angles at the base is  $30^\circ$ . Find the locus of the vertex.

12. Shew that the lines drawn from the angles of a triangle perpendicular to the opposite sides meet at a point; and that the locus of that point, for all equal triangles on the same base, is a parabola whose latus rectum is the altitude of the triangle.

## ANSWERS TO THE EXERCISES.

[B].

1.  $7x+8y=19$ .                      2.  $19x-11y=132$ .
3.  $3x=7y$ ; and  $3x-7y=11$ .      4.  $3x-5y=20$ .
5.  $109y+(448\pm 305\sqrt{3})x-5(1102\pm 305\sqrt{3})=0$ .
6. 20.      7. 35.      9.  $\frac{4}{3}$ .      10.  $4x+5y=0$ .      11.  $1383\frac{1}{2}$ .
12. 383.      13. Taking  $(x', y')$  for P, and  $(x'', y'')$  for Q,  
area  $=\frac{1}{2}(x'y'-x'y'')$ ; OP, OQ, PQ, = 13, 15,  $\sqrt{58}$ .
14. (2, 3), (5, 2), (-8, 28); area  $=32\frac{1}{2}$ .      15.  $y=\frac{7}{4}x+\frac{11}{4}$ ;  
DE is parallel to AC.      16.  $(a'-a)y-(b'-b)x-a'b$   
 $+ab'=0$ ; and  $(a'-a)y+(b'-b)x-a'b'+ab=0$ .
17. The eqn. to OR is  $x=0$ ; QR,  $4y-3x=56$ ;  
PR,  $5x+12y=168$ ; area of paral<sup>m</sup> = 168.
18.  $y=-3x+9$ ;  $3y=x+7$ ;  $y=-3x+14$ ;  $3y=x+12$ .
19.  $391x+187y=2184$ .      21.  $\tan CAB=m'$ ;  $\tan CBA$   
 $=-m$ ;  $\tan ACB=\frac{m-m'}{1+mm'}$ ;  $AB=\frac{c}{m}$ ;  
 $AC=\frac{c}{m-m'}\sqrt{1+m'^2}$ ;  $BC=\frac{cm'}{m(m-m')}\sqrt{1+m^2}$ .

[C].

1. The radius and the coordinates of the centre are, (i.) 7;  
3 and 4; (ii.) 3; -1 and -2; (iii.)  $4\sqrt{5}$ ; 3 and 9;  
(iv.)  $\sqrt{5}$ ;  $-\frac{1}{2}$  and 1.
2. (i.) (5, 3) and (-3, -5); (ii.)  $(2\frac{1}{2}, -5\frac{1}{2})$  and (-3, 5).
3. (i.) The line is a tangent to the circle at the point  
 $(\frac{3}{8}, -\frac{4}{8})$ ; (ii.) The line meets the circle at the point  
(-1, 0), which is an extremity of the diameter in the  
axis of  $x$ , and at the point  $(-\frac{9}{4}, -\frac{49}{4})$ .

4. (i.) Origin on the circumf.; points of intersection (24, 10), which is an extremity of the diam. through the origin, and (3.84, -5.12). (ii.) Points of intersection (6, -3½), (4, -5).

5.  $y - b = m(x - a) \pm c\sqrt{1 + m^2}$ . 6. 32.

7.  $\frac{1}{a^2} + \frac{1}{b^2} = \frac{1}{c^2}$ .

8.  $x = \frac{1}{2a}(a^2 + c^2 - c'^2)$ ;  $y = \pm \frac{1}{2a}\sqrt{(a + c + c')(a + c - c')(a - c + c')(c - a + c')}$ .

9. The centre is on the axis of  $x$  at  $\frac{m^2 + 1}{m^2 - 1}a$  from the

origin; the rad. =  $\frac{2m}{m^2 - 1}a$ ; coordinates of points of

intersection with the axis of  $x$  are  $x = \frac{m + 1}{m - 1}a$  and

$$x = \frac{m - 1}{m + 1}a.$$

11.  $x^2 + y^2 = 2\frac{1}{2}$ .

12.  $x^2 + y^2 - ax = c^2 - \frac{1}{2}a^2$ .

13.  $c = \pm \frac{1}{2}\sqrt{2(a - b)}$ . 14.  $x^2 + y^2 - \frac{2c}{1 + m^2}(x - my) = 0$ .

[D].

1. (i.)  $\frac{1}{2}$ ; (ii.)  $\frac{1}{2}\sqrt{10}$ ; (iii.)  $\frac{1}{2}\frac{2}{3}$ . 2.  $e = \frac{1}{2}\sqrt{2}$ .

3.  $1\frac{1}{4}$  and  $\frac{3}{4}\sqrt{15}$ . 4.  $y + ae^2 = \frac{x}{e}$ . 5. 21 and 15.

10.  $3y + \sqrt{5}x - 9a = 0$ ; the intercept on the axis of  $y$  is  $3a$ ; that on the axis of  $x$  is  $\frac{9a}{5}\sqrt{5}$ .

[E].

2.  $e = \frac{1}{2}\sqrt{10}$ ; lat. rect. =  $3a$ . 5.  $\frac{a^2}{n^2} - \frac{b^2}{m^2} = 1$ .

7. An hyperbola, if N be between A and B, and be not the middle point of AB. If N be on AB produced, the locus is an ellipse.

[F].

1. An angle whose tangent is  $\frac{1}{3}$ .
2.  $\frac{4}{3}a$ .
3.  $5a$ .
5.  $8a\sqrt{2}$ .
6. A parabola of which a tangent to the circle at A is the directrix, and the centre of the circle is the focus.

[G].

2.  $x'y' = \frac{1}{2}a^2$ .
3. Two parallels, each inclined to the axis of  $x$  at an angle whose tangent  $= -\frac{3}{4}$ , and whose distance from each other  $= \pm \frac{1}{10}\sqrt{13}$ .
4. Axes of the ellipse,  $2a = \frac{2}{3}\sqrt{6}$ ,  $2b = 2\sqrt{2}$ .
5. Two parallels whose eqns. are  $y = \frac{2}{3}x + \frac{2}{3}$  and  $y = \frac{2}{3}x + 1$ .
6. Axes of the hyperbola,  $2a = 2\sqrt{2}$ , and  $2b = 2$ .
7. A rectangular hyperbola, each semiaxis  $= 2^{\frac{1}{2}}$ .
8. A parabola; the coordinates  $1\frac{1}{2}\sqrt{2}$  and  $-\sqrt{2}$ .
9. A parabola; latus rectum  $= \sqrt{2}$ .
10. An ellipse; the coordinates of the centre  $\frac{1}{2}$  and  $-\frac{1}{2}$ , and the axes  $1\frac{1}{2}\sqrt{2} + \sqrt{2}$  and  $1\frac{1}{2}\sqrt{2} - \sqrt{2}$ .
11. An equilateral hyperbola whose axes are each  $= \frac{1}{2}\sqrt{2n}$ , and whose transverse axis makes with the base of the triangle an angle  $= 30^\circ$ .

LONDON

PRINTED BY SPOTTISWOODE AND CO.  
NEW-STREET SQUARE





## *Works by the same Author.*

---

### I

**MODERN ARITHMETIC.** A Treatise adapted for School Work and for Private Study. Containing numerous Improvements in aid of the Preparation of Candidates for Public Examinations.....12mo. [*Nearly ready.*]

### II

**EXERCISES in the FIRST FOUR RULES of ARITHMETIC,** constructed for the Application of New Artificial Tests, by which the Teacher may expeditiously ascertain the Correctness of the Results. Third Edition.....12mo. 6d.

### III

**SOLUTIONS of QUESTIONS in ARITHMETIC and BOOK-KEEPING** used in the Civil Service Examinations of 1882, and published in the Appendix to the Eighth Report of the Commissioners: with a SUPPLEMENT containing Examples in Account-States, &c. ....12mo. 1s. 6d.

### IV

**MERCANTILE EXERCISES in ADDITION of MONEY** and in the CALCULATION of PER-CENTAGES; including numerous Examples of a General and Simple Method of treating Per-Centage Problems, with Answers to the Exercises .....12mo. 1s. 3d.

The publication of this manual is the result of a conviction that, in a course of training for clerkships in Government offices, mercantile houses, &c. there is wanted a means of attaining a more thorough mastery of the general principles of per-centage calculation, and a more ready power of totalling long columns of money, than any of the published treatises on arithmetic are fitted

to impart. The instructions and exercises here supplied form a useful supplement to the ordinary arithmetical work in schools, besides affording help to candidates preparing for any of the various public examinations in which Addition of Money forms the subject of a special paper, and in which Problems in Percentages form a prominent feature of the miscellaneous paper.

### V

**PROGRESSIVE EXERCISES in BOOK-KEEPING by DOUBLE ENTRY,** including Account-States, Partnership Accounts, Private Journal, and Ledger, &c. ....12mo. 1s. 6d.  
'One of the most practical instructors in the art of *double entry*.....All schools should have it.'  
JOHN BULL.

### VI

**EXAMINATION-QUESTIONS in BOOK-KEEPING by DOUBLE ENTRY,** preceded by full Directions for the True Stating of Dr. and Cr.; with ANSWERS, providing the means of solving all the Problems and Exercises relating to Account-Books in the Civil Service and other Examination-Papers .....12mo. 2s. 6d.

'This volume is based on the examination papers published in the reports of the Civil Service Commissioners, which supply an excellent means of tuition and discipline. Some of these papers, however, although presenting in a small compass a great variety of transactions, and such as occur in actual business, still leave many who have gone through a course of instruction in book-keeping at a loss for the technical form and language usually adopted in practice. To afford

help in this difficulty, Mr. HUNTER has arranged some of the more useful of these papers, besides introducing some original questions and examples; and he has added answers to the whole, as a source of consultation in case of doubt and of correction in case of error. The work is adapted for use in schools generally, and provides the means of easily understanding and correctly treating the peculiarities of this department of mercantile affairs.'

MIDLAND COUNTIES PRESS.

## VII

**HUNTER'S EXAMINATION-QUESTIONS and DIRECTIONS**, as above, separated from the **ANSWERS**.....12mo. 1s.

## VIII

**RULED PAPER** for the various Forms of Account Books required in **HUNTER'S Examination-Questions in Book-keeping**, 5 sorts, price 1s. 6d. per Quire.

## IX

**EXAMINATION-QUESTIONS on COLENSO'S ELEMENTS of ALGEBRA, PART I.** .....12mo. 2s. 6d.

'THOSE teachers who use the **FIRST PART of COLENSO'S Algebra**, but who are unable themselves to prepare examination questions thereon, have to thank Mr **HUNTER** for doing this very thoroughly for them. He has appended some elementary exercises, suggested the re-

arrangement of several sets of exercises in **COLENSO**, and added a useful classification of problems in simple and quadratic equations, as well as examples of the application of algebra to geometrical constructions.' **MUSNUM.**

## X

**TREATISE on LOGARITHMS: With Copious Tables of Selected Logarithms; explaining simply the Nature and use of Logarithms and Logarithmic Tables, the Principles and Methods of their Construction, and their Application: With numerous Examples and Exercises** .....18mo. 1s. **KEY, 9d.**

'THIS treatise is likely to be useful. It has plenty of examples. **[ATHENZÆUM.**

'Its simplicity and clearness recommend this elementary treatise as a useful introduction to the study of logarithms generally, and especially as a preparation for more elaborate works on the subject.' **EDUCATIONAL TIMES.**

'MR. **HUNTER** in the present treatise explains in as simple a manner as pos-

sible the theory and practice of logarithms, a knowledge of which is now frequently required in candidates for public examination, in addition to being taught in the higher classes of schools. The little volume forms one of the well-known series of cheap school-books edited by the **Rev. G. E. GLEIG** which have rendered such valuable aid in the work of elementary education.'

**MIDLAND COUNTIES HERALD.**

## XI

**ELEMENTS of MENSURATION, simplified for the use of Beginners; with numerous Original Problems and Progressive Exercises. New Edition, Revised and Stereotyped** .....18mo. 9d.—**KEY, 9d.**

## XII

**ELEMENTS of PLANE TRIGONOMETRY, for Beginners; with numerous Problems, and Tables of all the Natural Sines, &c. required for the Solution of the Exercises** .....18mo. 1s.—**KEY, 9d.**

## XIII

**TEXT-BOOK of ENGLISH GRAMMAR: a Treatise on the Etymology and Syntax of the English Language; including Exercises in Parsing and Punctuation: an Etymological Vocabulary of Grammatical Terms; and a copious List of the principal Works on English Grammar. New Edition, and thoroughly Revised** .....12mo. 2s. 6d.

XIV

**EXERCISES in ENGLISH PARSING**, Progressively arranged and adapted to the Author's *Text-Book of English Grammar*; with Questions suggesting a Course of Oral Instruction for Junior Pupils. Tenth Edition ..... 12mo. 6d.

XV

**PARAPHRASING and ANALYSIS of SENTENCES**, simplified for the use of Schools, forming a Manual of Instruction and Exercise for the use of Students, Teachers, &c. .... 12mo. 1s. 3d.—KEY, 1s. 3d

XVI

**SCHOOL MANUAL of LETTER WRITING**: Containing numerous Models of Letters on Commercial and other subjects; with Exercises in Epistolary Composition, Rules of Punctuation, Explanations of Abbreviated Titles, Commercial Terms, &c. Second Edition ..... 12mo. 1s. 6d.

'This is one of the most serviceable books that have of late years been added to those in ordinary school use. Teachers have long experienced the want of such a book. The compilations called *Letter Writers* are, for the most part, such as might, without any loss to society in general, or to schools in particular, be consigned to everlasting oblivion. On the other hand, Mr. HUNTER'S *School*

*Manual of Letter Writing* is a work which should be introduced into every school where English composition is taught. The genuineness and modern style of the various commercial letters included in the present collection recommend themselves to the special attention of youths destined for mercantile life, as an excellent means of promoting in their minds an aptitude for the pursuits of business.'

PUPIL TEACHER.

XVII

**INTRODUCTION to the WRITING of PRECIS or DIGESTS**, as applicable to Narratives of Facts or Historical Events, Correspondence, Evidence, Official Documents, and General Composition: with numerous Examples and Exercises. Fourth Edition ..... 12mo. 2s.—KEY, 1s.

To this edition a new chapter has been added, to explain and exemplify that kind of précis or abstract which reduces a series of letters or other detached pieces to a continuous narrative: a

branch of précis-writing in which Candidates for the Civil Service Examinations are now generally required to possess readiness and proficiency.

XVIII

**JOHNSON'S RASSELAS**, with Introductory Remarks, Explanatory and Grammatical Annotations, Specimens of Interrogative Lessons, Answers to Examination Questions, &c. .... 12mo. 2s. 6d.

XIX

**BACON'S ADVANCEMENT of LEARNING**, Annotated Summary of Books I. and II. with numerous Extracts from the Work and Specimens of Examination-Questions ..... 12mo. 2s.

XX

**EXAMINATION-QUESTIONS on the FIRST TWO BOOKS of MILTON'S PARADISE LOST**, and on SHAKESPEARE'S *MERCHANT OF VENICE*, for the use of Middle-Class Candidates ..... 12mo. 1s.

XXI

**SELECT PLAYS of SHAKSPEARE**, with Explanatory and Illustrative Notes, Critical Remarks, and other Aids to a thorough understanding of each Drama. Edited for the use of Schools and Pupils preparing for Examination ..... 12mo. 2s. 6d. each Play.

HENRY VIII.  
JULIUS CÆSAR.

MERCHANT of VENICE.  
HAMLET.

THE TEMPEST.  
KING LEAR.

\* Certain departments in the Civil Service and the University Middle-class Examinations have of late given rise to a kind of school-book altogether new—prose translations of English poets. Thus many of the plays of SHAKSPEARE and the poems of MILTON, to insure their clearer comprehension—have been robbed of their metrical dress, and have been introduced to us in the humbler garb of prose—the original and the paraphrase

being placed side by side. The reader is thus shown the peculiar idioms of his Author; he gradually contracts a habit of analysing what he reads; and in the case of other works, the structure of whose sentences is not that of the present day, of being dissatisfied, unless he is able at each step to see clearly the force of every particle and every turn of expression.  
READER.

XXII

**BOOKS I. and II. of MILTON'S PARADISE LOST:**

with a Prose Translation or Paraphrase, the Parsing of the more Difficult Words, Specimens of Analysis, and numerous Illustrative Notes, for the use of Candidates..... 12mo. price 1s. 6d. each Book.

\* The announcement of the University of Oxford, that the First Book of *Paradise Lost* would form a subject on which junior candidates for the Middle-Class Examinations must be prepared, suggested the present attempt to provide a means of guidance and instruction for the special preparation of such candidates, and at the same time a schoolbook of general utility. To this explanation of the origin and object of the book before us, we have only to add our sincere wish, that it may have a circulation commensurate with its merits. In that case, the finest epic poem in the English language, which has won for its author an imperishable renown, will be far more extensively read and appreciated than it has yet been, even among those who have received what is generally understood to be a good education. The chief reason for the existence of this comparative ignorance of MILTON'S immortal poem is

clearly pointed out by Mr. HUNTER, when he says: "It is a very profound work, abounding in unexploited, idealising thought, copiously characterised by imitations of the transpositive, elliptical, and idiomatic structure of the ancient languages of Greece and Rome, and affluent in allusions to ancient history, fable, and romance. It is, therefore, a hard book for the general reader, and a study for the classical scholar." The difficulties indicated in this passage will be found to be either greatly diminished, or entirely removed in the volume before us, thus rendering familiar and intelligent acquaintances with the first book at least of *Paradise Lost* a work of comparative ease. Whoever advances thus far will not stand in need of a stimulus to urge him to master the remainder of the poem.

BELL'S WEEKLY MESSENGER.

XXIII

**MILTON'S COMUS, L'ALLEGRO, and IL PENSE-**

*EOSO*, with numerous Notes, critical, illustrative, and explanatory: adapted for use in Training Colleges and Schools ..... 12mo. 1s. 6d.

\* The notes are short, terse, and quite to the purpose. GUARDIAN.

\* An edition carefully illustrated with notes, and well suited for training colleges and schools. JOHN BULL.

\* Mr. HUNTER is among the best of the numerous editors of our classical writers whom the University Middle Class examinations have called out. In place of criticism of his own he gives by way of introduction copious extracts from the criticisms of WARTON, JOHNSON, HALLAM, MACAULAY, and JAMES MONTGOMERY; and his notes are terse and to the point, while

they give all the information even school-boys can require. The plan of explaining unusual words or constructions by parallel passages from other poets is, we think, far better than a laboured explanation. SPECTATOR.

\* The notes to this edition, which are very comprehensive, are not restricted to mere exegesis; they explain difficult and obscure passages by parallel constructions from other poets. The critical extracts prefixed convey a just and clear appreciation of the peculiarities of MILTON'S style.  
SUN.

London LONGMANS, GREEN, and CO. Paternoster Row.





## REDUCTION IN THE PRICE OF LAURIE'S GRADUATED ENGLISH READING BOOKS.

‘Mr. LAURIE’s excellent *Graduated Series of Reading-Lesson Books* has met  
with great and well-deserved success.’ ATHENÆUM.

Complete in 5 Books, with New PRIMER, price 8s. 1d.

# THE GRADUATED SERIES OF READING-LESSON BOOKS

WITH EXPLANATORY NOTES, ADAPTED  
AS A PROGRESSIVE COURSE OF READING, FOR ALL CLASSES OF  
ENGLISH SCHOOLS, INSTITUTIONS, AND FAMILIES.

EDITED BY J. S. LAURIE,

LATELY ONE OF HER MAJESTY’S INSPECTORS OF SCHOOLS.

Messrs. LONGMANS and Co. beg leave to announce the reissue of Mr. LAURIE’s *Graduated Series of Reading Books* at considerably reduced prices. They also have to announce that to render them at once more complete, and better suited to the present requirements of Inspected and other Schools, a new Primer suited to Standard I. of the Revised Code has been added to them by Dr. J. D. MORELL, one of H.M. Inspectors of Schools; and a careful selection of poetry appended to Books III. and IV. All the

books have been several times re-printed, and now and revised editions are now ready.

The *Graduated Series* in its present form not only comprehends a far greater quantity of matter than any of the other Standard Reading Books—but presents a variety of Readings on Historical, Geographical, Scientific, and other Miscellaneous Subjects; which have been acknowledged to be unequalled by any similar publication. The prices at which the Series is now offered are the following:—

Elementary Reading Book, or Primer, by Dr. J. D. MORELL, suited to Standard I. pp. 60, price 2d. sewed, or 4d. cloth...	0s. 2d.
Book I. suited to Standard II. pp. 192.....	0s. 8d.
Book II. suited to Standards III. and IV. pp. 254.....	1s. 3d.
Book III. suited to Standards IV. and V. pp. 312.....	1s. 6d.
Book IV. suited to Standard VI. pp. 440.....	2s. 0d.
Book V. comprehending Readings in the best English Literature of the present day, pp. 496.....	2s. 6d.

Taking into account the amount of matter here presented, and the fact that ample materials for Standards III. IV. and V. are contained in two of the above volumes (so that it is not

necessary to purchase a new book for every separate Standard), the *Graduated Series* are amongst the cheapest Reading Books ever offered to the public.

‘We have had occasion to speak in terms of the highest approbation of two or three volumes of the *Graduated Series of Reading-Lesson Books*. The fifth and last book of the series, now before us, shows how excellently the plan adopted in the arrangement of the books preceding it has been calculated to lead the pupil onwards to a full appreciation of these complementary selections from the higher branches of literature. Steadily, yet surely, the pupil has been led onward from the simplest narratives, the plainest told tales, and the most lucid descriptions of common objects, to a point in which exactly similar departments of literature are represented by specimens from the works of some of the finest writers of the last and present century.’ CHURCHMAN.

London: LONGMANS, GREEN, and CO. Paternoster Row.



